

Mouse anti-Human Vimentin Monoclonal Antibody | anti-VIM antibody

Vimentin (Mesenchymal Cell Marker)

CATALOG # AAA215254	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	----------------------	----------------------------	---

PRODUCT SYNONYMS

Vimentin; Monoclonal Antibody; Vimentin (Mesenchymal Cell Marker); VIM; anti-VIM antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	V9
Specificity	This MAb reacts with a 58kDa protein identified as vimentin. It shows no cross-reaction with other closely related intermediate filament proteins (IFP's) such as desmin, keratin, neurofilament, and glial fibrillary acid protein. Anti-vimentin alone is of limited value as a tool; however, when used in panels with other antibodies, it is useful for the sub-classification of a given tumor. Expression of vimentin, when used in conjunction with anti-keratin, is helpful when distinguishing melanomas from undifferentiated carcinomas and large cell lymphomas. All melanomas and Schwannomas react strongly with anti-vimentin. It labels a variety of mesenchymal cells, including melanocytes, lymphocytes, endothelial cells, and fibroblasts. Non-reactivity of anti-vimentin is often considered more useful than its positive reactivity, since there are a few tumors that do not contain vimentin, e.g. hepatoma and seminoma. Anti-vimentin is also useful as a tissue process control reagent.
Purity/Purification	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml
Form/Format	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	466
Applicable Applications for anti-VIM antibody	IHC (Immunohistochemistry), WB (Western Blot)
Positive Control	U-87, Raji, Jurkat or HeLa cells. Sarcomas or Melanomas.
Immunogen	Porcine Lens
Cellular Localization	Cytoplasmic
Hu-chromosome Location	10p12.33
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C. Antibody without azide: store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



WB (Western Blot)
Western Blot Analysis of U-87 cell lysate using Vimentin Mouse Monoclonal Antibody (V9).



SDS-PAGE
SDS-PAGE Analysis Purified Vimentin Mouse Monoclonal Antibody (V9). Confirmation of Integrity and Purity of Antibody.



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Uterine Carcinoma stained with Vimentin Mouse Monoclonal Antibody (V9).

ADDITIONAL INFORMATION

References	Osborn M et. al. European Journal of Cell Biology. 1984; 34:137-143
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	62414289
NCBI GeneID	7431
NCBI Accession #	NP_003371.2
NCBI GenBank Nucleotide #	NP_003371.2
UniProt Accession #	P08670
Molecular Weight	57-60kDa
NCBI Official Full Name	vimentin
NCBI Official Synonym Full Names	vimentin
NCBI Official Symbol	VIM
NCBI Protein Information	vimentin
UniProt Protein Name	Vimentin
UniProt Gene Name	VIM
UniProt Entry Name	VIME_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/vimentin-antibody-clone-v9-immunohistochemistry-western-blot-elisa/215254>