

Mouse anti-Human CD8A Monoclonal Antibody | anti-CD8A antibody

CD8A (Cytotoxic- & Suppressor T-Cell Marker)

CATALOG # AAA215289	HOST Mouse	REACTIVITY Human	APPLICATIONS Flow Cytometry, Functional Assay
-------------------------------	----------------------	----------------------------	---

PRODUCT SYNONYMS

CD8A; Monoclonal Antibody; CD8A (Cytotoxic- & Suppressor T-Cell Marker); CD8 antigen; alpha polypeptide (p32); T8/Leu-2 T-lymphocyte differentiation antigen; Ly3; LYT3; MAL; T-cell surface glycoprotein CD8 alpha chain; anti-CD8A antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG2a, kappa
Clone Number	UCHT4
Specificity	CD8 is a cell surface receptor expressed either as a heterodimer with the CD8 beta chain (CD8 alpha/beta) or as a homodimer (CD8 alpha/alpha). A majority of thymocytes and a subpopulation of mature T cells and NK cells express CD8a. CD8 binds to MHC class 1 and through its association with protein tyrosine kinase p56lck plays a role in T cell development and activation of mature T cells. For mature T-cells, CD4 and CD8 are mutually exclusive, so anti-CD8, generally used in conjunction with anti-CD4. It is a useful marker for distinguishing helper/inducer T-lymphocytes, and most peripheral T-cell lymphomas are CD4+/CD8-. Anaplastic large cell lymphoma is usually CD4+ and CD8-, and in T-lymphoblastic lymphoma/leukemia, CD4 and CD8 are often co-expressed. CD8 is also found in littoral cell angioma of the spleen. This MAb binds to the CD8 alpha chain, at an epitope distinct from that recognized by UCHT14.
Purity/Purification	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml
Form/Format	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	235
Applicable Applications for anti-CD8A antibody	FCM/FACS (Flow Cytometry)
Positive Control	Jurkat, HuT78 or hPBL. Tonsil.
Immunogen	Thymocytes followed by Sezary T cells.
Cellular Localization	Cell Surface
Hu-chromosome Location	2p11.2
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C. Antibody without azide: store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



FCM/FACS (Flow Cytometry)
Flow Cytometric Analysis of Jurkat cells. CD8 Mouse Monoclonal Antibody (UCHT4) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).



SDS-PAGE
SDS-PAGE Analysis Purified CD8 Mouse Monoclonal Antibody (UCHT4). Confirmation of Purity and Integrity of Antibody.

ADDITIONAL INFORMATION

References	Mason DY, et. al. Journal of Clinical Pathology, 1992, 45(12):1084-8.
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	225007536
NCBI GeneID	925
NCBI Accession #	NP_001139345.1
NCBI GenBank Nucleotide #	NP_001139345.1
UniProt Accession #	P01732
Molecular Weight	32kDa
NCBI Official Full Name	T-cell surface glycoprotein CD8 alpha chain isoform 1
NCBI Official Synonym Full Names	CD8a molecule
NCBI Official Symbol	CD8A
NCBI Official Synonym Symbols	CD8; p32; Leu2
NCBI Protein Information	T-cell surface glycoprotein CD8 alpha chain
UniProt Protein Name	T-cell surface glycoprotein CD8 alpha chain
UniProt Gene Name	CD8A
UniProt Synonym Gene Names	MAL
UniProt Entry Name	CD8A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd8a-antibody-clone-ucht4-flow-cytometry-elisa/215289>