

Mouse anti-Human CD22/BL-CAM Monoclonal Antibody | anti-CD22 antibody

CD22/BL-CAM (B-Cell Marker)


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA215297	Mouse	Human	Immunofluorescence, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS


CD22/BL-CAM; Monoclonal Antibody; CD22/BL-CAM (B-Cell Marker); B-lymphocyte cell adhesion molecule (BL-CAM); B-cell receptor CD22; CD22; Lectin 2; Lyb8; Sialic acid-binding Ig-like lectin 2 (Siglec-2); SIGLEC2; T-cell surface antigen Leu-14; anti-CD22 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	RFB4
Specificity	Recognizes a protein of 130-140kDa, identified as CD22 (also known as BL-CAM). CD22 expression is restricted to normal and neoplastic B cells and is absent from other haemopoietic cell types. In B-cell ontogeny, CD22 is first expressed in the cytoplasm of pro-B and pre-B cells, and on the surface as B cells mature to become IgD+. It is not expressed by plasma cells, CD22 is found highly expressed in follicular mantle and marginal zone B-cells, and while germinal center B-cells are relatively weak. CD22 is a member of the immunoglobulin superfamily and serves as an adhesion receptor for sialic acid-bearing ligands expressed on erythrocytes and all leukocyte classes. It also associates with tyrosine kinases and play a role in signal transduction and B-cell activation.
Purity/Purification	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml
Form/Format	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	759
Applicable Applications for anti-CD22 antibody	IF (Immunofluorescence), FCM/FACS (Flow Cytometry), ELISA
Positive Control	Ramos, Raji or Daudi cells. Human Tonsil, Lymph Node or Spleen.
Immunogen	Human tonsil lymphocytes.
Cellular Localization	Cell Surface
Hu-chromosome Location	19q13.1
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C. Antibody without azide: store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.



FCM/FACS (Flow Cytometry)
Flow Cytometric Analysis of Ramos cells using CD22 Mouse Monoclonal Antibody (RFB4) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype Control (Red).



SDS-PAGE
SDS-PAGE Analysis Purified CD22 Mouse Monoclonal Antibody (RFB4). Confirmation of Integrity and Purity of Antibody.

ADDITIONAL INFORMATION

References	Sato S et al. Semin Immunol 1998, 10:287-297
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	297374825
NCBI GeneID	933
NCBI Accession #	NP_001172028.1
NCBI GenBank Nucleotide #	NP_001172028.1
UniProt Accession #	P20273
Molecular Weight	130-140kDa
NCBI Official Full Name	B-cell receptor CD22 isoform 2
NCBI Official Synonym Full Names	CD22 molecule
NCBI Official Symbol	CD22
NCBI Official Synonym Symbols	SIGLEC2; SIGLEC-2
NCBI Protein Information	B-cell receptor CD22
UniProt Protein Name	B-cell receptor CD22
UniProt Gene Name	CD22
UniProt Synonym Gene Names	SIGLEC2; BL-CAM; Siglec-2
UniProt Entry Name	CD22_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.