

Mouse anti-Human CD23 Monoclonal Antibody | anti-FCER2 antibody

CD23 (Fc Epsilon RI) Mouse Monoclonal Antibody [Clone FCER2/6887]

CATALOG # AAA215643	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

CD23; Monoclonal Antibody; CD23 (Fc Epsilon RI) Mouse Monoclonal Antibody [Clone FCER2/6887]; CD23 (Fc Epsilon RI); BLAST-2; C-type lectin domain family 4; member J; Fc fragment of IgE low affinity II receptor; Fc-epsilon-RII; FCER2; FCER2A; FceRII; IgE-binding factor (IGEBF); Immunoglobulin E receptor; low affinity II; Lymphocyte IgE receptor; anti-FCER2 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse/IgG2b, kappa
Clone Number	FCER2/6887
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Concentration	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml (varies by lot)
Applicable Applications for anti-FCER2 antibody	IHC (Immunohistochemistry)
Immunogen	Recombinant fragment (around aa 48-321) of human FCER2/CD23 protein (exact sequence is proprietary)
Cellular Localization	Cell Surface
Positive Control	Human lymph node or tonsil (IHC).
Hu-chromosome Location	19p13.3
Unigene	465778
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C Antibody without azide: store at -20 to -80 degree C Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



SDS-PAGE

SDS-PAGE Analysis Purified CD23-Monospecific Mouse Monoclonal Antibody (FCER2/6887).
Confirmation of Purity and Integrity of Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-FCER2 antibody	CD23 (FCE2) is a type II integral membrane glycoprotein that is expressed on mature B cells, monocytes, eosinophils, platelets and dendritic cells. CD23 is a low affinity IgE receptor that mediates IgE-dependent cytotoxicity and phagocytosis by macrophages and eosinophils. CD23 associates as an oligomer where cooperative binding of at least two lectin domains is required for high affinity IgE binding to CD23. It may play a role in antigen presentation by B cells by interacting with CD40. CD23 has been shown to be associated with the Fyn tyrosine kinase. The truncated molecule can be secreted, then function as a potent mitogenic growth factor. CD23 is expressed on a subpopulation of peripheral blood cells, B-lymphocytes and on EBV transformed B lymphoblastoid cell lines. CD23 is also detected in neoplastic cells from cases of B cell chronic lymphocytic leukemia and some cases on centroblastic/centrocytic lymphoma.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	119862
NCBI GeneID	2208
UniProt Accession #	P06734
Molecular Weight	321
NCBI Official Full Name	Low affinity immunoglobulin epsilon Fc receptor
NCBI Official Synonym Full Names	Fc fragment of IgE, low affinity II, receptor for (CD23)
NCBI Official Symbol	FCER2
NCBI Official Synonym Symbols	CD23; FCE2; CD23A; IGEBF; CLEC4J; BLAST-2
NCBI Protein Information	low affinity immunoglobulin epsilon Fc receptor; CD23 antigen; fc-epsilon-RII; lymphocyte IgE receptor; immunoglobulin E-binding factor; C-type lectin domain family 4, member J
UniProt Protein Name	Low affinity immunoglobulin epsilon Fc receptor
UniProt Gene Name	FCER2
UniProt Synonym Gene Names	CD23A; CLEC4J; FCE2; IGEBF
UniProt Entry Name	FCER2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd23-antibody-clone-fcer2-6887-immunohistochemistry-elisa/215643>