

Mouse anti-Human GC Vitamin D Binding Protein Monoclonal Antibody | anti-GC antibody

GC Vitamin D Binding Protein Mouse Monoclonal Antibody [Clone VDBP/4481]

CATALOG # AAA215675	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

GC Vitamin D Binding Protein; Monoclonal Antibody; GC Vitamin D Binding Protein Mouse Monoclonal Antibody [Clone VDBP/4481]; DBP/GC; Gc globulin; GRD3; Group specific component vitamin D binding protein; hDBP; VDB; VDBG; VDBP; Vitamin D binding alpha globulin; Vitamin D-binding protein; anti-GC antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse/IgG2b, kappa
Clone Number	VDBP/4481
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Concentration	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml (varies by lot)
Applicable Applications for anti-GC antibody	IHC (Immunohistochemistry)
Immunogen	Recombinant fragment (around aa 35-175) of human GC protein (exact sequence is proprietary)
Cellular Localization	Secreted
Positive Control	Human liver, kidney or pancreas.
Hu-chromosome Location	4q13.3
Unigene	418497
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C Antibody without azide: store at -20 to -80 degree C Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



SDS-PAGE
SDS-PAGE Analysis Purified GC Mouse Monoclonal Antibody (VDBP/4481). Confirmation of Purity and Integrity of Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-GC antibody	Vitamin D-binding protein (DBP) is a multi-functional serum protein that binds to the plasma membranes of numerous cell types and mediates a variety of cellular functions. The locus of the DBP protein (also known as group-specific component protein or GC) is located at human chromosome 4q13. 3. DBP functions in organ-specific transportation of vitamin D and its metabolites to the various target organs of the vitamin D endocrine system. In addition, DBP has immunomodulatory properties and is able to bind to the surface of leukocytes. DBP binds to the plasma membrane through a chondroitin sulfate proteoglycan. DBP serves as a co-chemotactic factor for C5a to enhance the chemotactic activity of C5a. DBP can also bind to globular Actin with high affinity and is involved in the clearance of Actin from the blood. DBP plays an important role in osteoclast differentiation. The diverse cellular functions of DBP require its cell surface binding ability to mediate different biological processes.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	324021745
NCBI GeneID	2638
NCBI Accession #	NP_001191236.1
NCBI GenBank Nucleotide #	NM_001204307.1
UniProt Accession #	P02774; Q16309; Q16310; Q53F31; Q6GTG1; B4DPP2; D6RAK8
Molecular Weight	52,964 Da
NCBI Official Full Name	vitamin D-binding protein isoform 3
NCBI Official Synonym Full Names	group-specific component (vitamin D binding protein)
NCBI Official Symbol	GC
NCBI Official Synonym Symbols	DBP; GRD3; VDBG; VDBP; DBP/GC
NCBI Protein Information	vitamin D-binding protein; VDB; gc-globulin; vitamin D-binding alpha-globulin
UniProt Protein Name	Vitamin D-binding protein
UniProt Gene Name	GC
UniProt Synonym Gene Names	DBP; VDB
UniProt Entry Name	VTDB_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/gc-vitamin-d-binding-protein-antibody-clone-vdbp-4481-immunohistochemistry-elisa/215675>