

Mouse anti-Human IRF3 Monoclonal Antibody | anti-IRF3 antibody

IRF3 Mouse Monoclonal Antibody [Clone PCRP-IRF3-1D11]

CATALOG # AAA215751	HOST Mouse	REACTIVITY Human	APPLICATIONS Western Blot, Immunofluorescence
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

IRF3; Monoclonal Antibody; IRF3 Mouse Monoclonal Antibody [Clone PCRP-IRF3-1D11]; IIAE7; MGC94729; Interferon regulatory factor 3 (IRF3); anti-IRF3 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse/IgG1, kappa
Clone Number	PCRP-IRF3-1D11
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Concentration	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml (varies by lot)
Applicable Applications for anti-IRF3 antibody	WB (Western Blot), IF (Immunofluorescence)
Immunogen	Recombinant full-length human IRF3 protein
Cellular Localization	Cytoplasm, Nucleus. Shuttles between cytoplasmic and nuclear compartments, with export being the prevailing effect. When activated, IRF3 interaction with CREBBP prevents its export to the cytoplasm
Positive Control	HeLa, U87 or Ramos cells.
Hu-chromosome Location	19q13.33
Unigene	289052
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C Antibody without azide: store at -20 to -80 degree C Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



SDS-PAGE
SDS-PAGE Analysis Purified IRF3 Mouse Monoclonal Antibody (PCRP-IRF3-1D11).
Confirmation of Purity and Integrity of Antibody.

ADDITIONAL INFORMATION

Related Product Information for anti-IRF3 antibody	Interferon regulatory factor-1 (IRF-1) and IRF-2 have been identified as novel DNA-binding factors that function as regulators of both type I interferon (interferon-alpha and beta) and interferon-inducible genes. The two factors are structurally related, particularly in their N-terminal regions, which confer DNA binding specificity. In addition, both bind to the same sequence within the promoters of interferon-alpha and interferon-beta genes. IRF-1 functions as an activator of interferon transcription, while IRF-2 binds to the same cis elements and represses IRF-1 action. IRF-1 and IRF-2 have been reported to act in a mutually antagonistic manner in regulating cell growth; overexpression of the repressor IRF-2 leads to cell transformation while concomitant overexpression of IRF-1 causes reversion. IRF-1 and IRF-2 are members of a larger family of DNA binding proteins that includes IRF-3, IRF-4, IRF-5, IRF-6, IRF-7, ISGF-3 gamma p48 and IFN consensus sequence-binding protein (ICSBP).
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	308199450
NCBI GeneID	3661
NCBI Accession #	NP_001184053.1
NCBI GenBank Nucleotide #	NM_001197124.1
UniProt Accession #	Q14653; Q5FBY1; A8K7L2
Molecular Weight	47,219 Da
NCBI Official Full Name	interferon regulatory factor 3 isoform 4
NCBI Official Synonym Full Names	interferon regulatory factor 3
NCBI Official Symbol	IRF3
NCBI Protein Information	interferon regulatory factor 3
UniProt Protein Name	Interferon regulatory factor 3
UniProt Gene Name	IRF3
UniProt Synonym Gene Names	IRF-3
UniProt Entry Name	IRF3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

