

Mouse anti-Human Cytokeratin 8 (KRT8) Monoclonal Antibody | anti-KRT8 antibody

Cytokeratin 8 (KRT8) Recombinant Mouse Monoclonal Antibody [Clone rKRT8/4209]

CATALOG # AAA215764	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

Cytokeratin 8 (KRT8); Monoclonal Antibody; Cytokeratin 8 (KRT8) Recombinant Mouse Monoclonal Antibody [Clone rKRT8/4209]; CARD2; CK8; CYK8; CYKER; Cytokeratin Endo A; DreK8; EndoA; K2C8; K8; Keratin 8; Krt 2. 8; KRT8; Type-II Keratin Kb8; anti-KRT8 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse/IgG2a, kappa
Clone Number	rKRT8/4209
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0. 05% BSA & 0. 05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Concentration	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml (varies by lot)
Applicable Applications for anti-KRT8 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Antibody Type	Recombinant Antibody
Immunogen	Recombinant full-length human KRT8 protein
Cellular Localization	Cytoplasmic
Positive Control	HCT116, MCF-7 or A431 cells. Human skin, colon, lung or breast carcinoma.
Hu-chromosome Location	12q13.13
Unigene	533782 & 708445
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C Antibody without azide: store at -20 to -80 degree C Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human colon stained with Cytokeratin 8 Recombinant Mouse Monoclonal Antibody (rKRT8/4209).

ADDITIONAL INFORMATION

Related Product Information for anti-KRT8 antibody	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenocarcinomas and ductal carcinomas. It is absent in squamous cell carcinomas. Hepatocellular carcinomas are defined by the use of antibodies that recognize only cytokeratin 8 and 18. CK8 exists on several types of normal and neoplastic epithelia, including many ductal and glandular epithelia such as colon, stomach, small intestine, trachea, and esophagus as well as in transitional epithelium. Anti-CK8 does not react with skeletal muscle or nerve cells. Epithelioid sarcoma, chordoma, and adamantinoma show strong positivity corresponding to that of simple epithelia (with antibodies against CK8, CK18 and CK19). Anti-CK8 is useful for the differentiation of lobular (ring-like, perinuclear) from ductal (peripheral-predominant) carcinoma of the breast.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	372466577
NCBI GeneID	3856
NCBI Accession #	NP_001243222.1
NCBI GenBank Nucleotide #	NM_001256293.1
UniProt Accession #	P05787; Q14099; Q14716; Q14717; Q53GJ0; Q6DHW5; Q6GMY0; Q6P4C7; Q96J60; A8K4H3; B0AZN5; F8VXB4
Molecular Weight	53,704 Da
NCBI Official Full Name	keratin, type II cytoskeletal 8 isoform 2
NCBI Official Synonym Full Names	keratin 8
NCBI Official Symbol	KRT8
NCBI Official Synonym Symbols	K8; KO; CK8; CK-8; CYK8; K2C8; CARD2
NCBI Protein Information	keratin, type II cytoskeletal 8; cytokeratin 8; cytokeratin-8; type-II keratin Kb8
UniProt Protein Name	Keratin, type II cytoskeletal 8
UniProt Gene Name	KRT8
UniProt Synonym Gene Names	CYK8; CK-8; K8
UniProt Entry Name	K2C8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

