

Rabbit anti-Human TACSTD2/TROP2 Monoclonal Antibody | anti-TACSTD2 antibody

TACSTD2/TROP2 (Epithelial Marker) Recombinant Rabbit Monoclonal Antibody [Clone TACSTD2/6396R]

CATALOG # AAA215780	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

TACSTD2/TROP2; Monoclonal Antibody; TACSTD2/TROP2 (Epithelial Marker) Recombinant Rabbit Monoclonal Antibody [Clone TACSTD2/6396R]; TACSTD2/TROP2 (Epithelial Marker); Cell surface glycoprotein Trop-2; Membrane Component Chromosome 1; Surface Marker 1 (M1S1); Pancreatic Carcinoma Marker Protein GA733-1; TROP2; Tumor-Associated Calcium Signal Transducer 2 (TACSTD2); anti-TACSTD2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Monoclonal
Isotype	Rabbit/IgG
Clone Number	TACSTD2/6396R
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Concentration	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml (varies by lot)
Applicable Applications for anti-TACSTD2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Antibody Type	Recombinant Antibody
Immunogen	Recombinant fragment of human TACSTD2 protein (around aa 31-274) (exact sequence is proprietary)
Cellular Localization	Cell Surface
Positive Control	HT29 cells. Human breast or colon carcinoma.
Hu-chromosome Location	1p32.1
Unigene	23582
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C Antibody without azide: store at -20 to -80 degree C Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)

Formalin-fixed, paraffin-embedded human thyroid carcinoma stained with TACSTD2 Recombinant Rabbit Monoclonal Antibody (TACSTD2/6396R).

ADDITIONAL INFORMATION

Related Product Information for anti-TACSTD2 antibody	TACSTD2 is a cell surface glycoprotein receptor. It is a single pass type I membrane protein containing one thyroglobulin type-1 domain, an epidermal growth factor-like repeat, a phosphatidylinositol binding site and tyrosine phosphorylation sites near the C-terminus. It plays a role in transducing intracellular calcium signals. It is expressed in trophoblast cells, cornea and multi-stratified epithelia. It is also highly expressed in several types of tumors and is involved in regulating the growth of carcinoma cells.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	166795236
NCBI GeneID	4070
NCBI Accession #	NP_002344.2
NCBI GenBank Nucleotide #	NM_002353.2
UniProt Accession #	P09758; Q15658; Q6FG48; Q7Z7Q4; Q96QD2
Molecular Weight	35,709 Da
NCBI Official Full Name	tumor-associated calcium signal transducer 2
NCBI Official Synonym Full Names	tumor-associated calcium signal transducer 2
NCBI Official Symbol	TACSTD2
NCBI Official Synonym Symbols	EGP1; GP50; M1S1; EGP-1; TROP2; GA7331; GA733-1
NCBI Protein Information	tumor-associated calcium signal transducer 2; epithelial glycoprotein-1; cell surface glycoprotein TROP2; cell surface glycoprotein Trop-2; pancreatic carcinoma marker protein GA7331; pancreatic carcinoma marker protein GA733-1; gastrointestinal tumor-ass
UniProt Protein Name	Tumor-associated calcium signal transducer 2
UniProt Gene Name	TACSTD2
UniProt Synonym Gene Names	GA733-1; M1S1; TROP2
UniProt Entry Name	TACD2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/tacstd2-trop2-antibody-clone-tacstd2-6396r-immunohistochemistry-western-blot-elisa/215780>