

Mouse anti-Human RBP4/Retinol Binding Protein 4 Monoclonal Antibody | anti-RBP4 antibody

RBP4/Retinol Binding Protein 4 Mouse Monoclonal Antibody [Clone RBP4/4048]

CATALOG # AAA215884	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

RBP4/Retinol Binding Protein 4; Monoclonal Antibody; RBP4/Retinol Binding Protein 4 Mouse Monoclonal Antibody [Clone RBP4/4048]; Plasma retinol binding protein 4 (PRBP); RDCCAS; Retinol binding protein 4 interstitial; anti-RBP4 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse/IgG2b, kappa
Clone Number	RBP4/4048
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Concentration	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml (varies by lot)
Applicable Applications for anti-RBP4 antibody	IHC (Immunohistochemistry)
Immunogen	Recombinant fragment (around aa29-148) of human RBP4 protein (exact sequence is proprietary)
Cellular Localization	Secreted
Positive Control	HeP-G2 cells. Human kidney, pancreas or liver tissues.
Hu-chromosome Location	10q23.33
Unigene	50223
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C Antibody without azide: store at -20 to -80 degree C Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human kidney stained with RBP4 Mouse Monoclonal Antibody (RBP4/4048).

ADDITIONAL INFORMATION

Related Product Information for anti-RBP4 antibody	Retinol (Vitamin A) is transported in the blood bound to its carrier protein, retinol-binding protein (RBP), also designated plasma retinol-binding protein (PRBP) or RBP4. A member of the lipocalin family, RBP conveys retinol from stores in the liver to peripheral tissues. In plasma, RBP binds transthyretin (TTR, formerly called prealbumin) to prevent glomerular filtration of low molecular weight RBP in the kidneys. The stability of this complex holds test importance because the molar ratio of RBP:TTR provides an indirect way to indicate marginal Vitamin A deficiency. Vitamin A deficiency blocks the secretion of RBP, resulting in defective delivery and supply to epidermal cells. Originally identified solely as a transporter protein, recent studies correlating increased levels of RBP expression in adipose tissue with Insulin resistance have generated research into the possible roles the protein may play in the pathogenesis of type 2 diabetes and obesity.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	55743122
NCBI GeneID	5950
NCBI Accession #	NP_006735.2
NCBI GenBank Nucleotide #	NM_006744.3
UniProt Accession #	P02753; O43478; O43479; Q5VY24; Q8WWA3; Q9P178; D3DR38
Molecular Weight	23,010 Da
NCBI Official Full Name	retinol-binding protein 4
NCBI Official Synonym Full Names	retinol binding protein 4, plasma
NCBI Official Symbol	RBP4
NCBI Official Synonym Symbols	RDCCAS
NCBI Protein Information	retinol-binding protein 4; RBP; PRBP; plasma retinol-binding protein; retinol-binding protein 4, interstitial
UniProt Protein Name	Retinol-binding protein 4
UniProt Gene Name	RBP4
UniProt Synonym Gene Names	PRBP; RBP
UniProt Entry Name	RET4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

