

Mouse anti-Human Replication Protein A2 (RPA2) Monoclonal Antibody | anti-RPA2 antibody

Replication Protein A2 (RPA2) Mouse Monoclonal Antibody [Clone RPA2/4774]

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA215897	Mouse	Human	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Replication Protein A2 (RPA2); Monoclonal Antibody; Replication Protein A2 (RPA2) Mouse Monoclonal Antibody [Clone RPA2/4774]; p32; p34; REPA2; Replication protein A 32kDa subunit; Replication protein A 34kDa subunit; Replication protein A2; RF-A protein 2; Rf-A2; RFA; RP-A p32; RP-A p34; RP21C; RPA32; RPA34; RpLP1; RpP2; anti-RPA2 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse/IgG1, kappa
Clone Number	RPA2/4774
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Concentration	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml (varies by lot)
Applicable Applications for anti-RPA2 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant human full-length Replication Protein A2 (RPA2) protein
Cellular Localization	Nuclear
Positive Control	MCF-7, HeLa or HepG2 cells. Human tonsil (IHC).
Hu-chromosome Location	1p35.3
Unigene	703070 & 79411
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C Antibody without azide: store at -20 to -80 degree C Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human colon stained with RPA2 Mouse Monoclonal Antibody (RPA2/4774).

ADDITIONAL INFORMATION

Related Product Information for anti-RPA2 antibody	Recognizes a protein of 34kDa, identified as a subunit of Replication Protein A (RPA) (also known as human single-stranded DNA binding protein, or HSSB). RPA from human cells is a stable heterotrimer of 70kDa, 32-34kDa, and 11-14kDa subunits (RPA70, RPA32, and RPA14 respectively). It is involved in DNA replication, repair, and recombination. RPA is required for the SV40 large tumor antigen-catalyzed unwinding of SV40 DNA and stimulates DNA polymerase (pol) alpha and delta. RPA34 is phosphorylated at the G1/S boundary of the cell cycle or upon exposure of cells to DNA damage-inducing agents including ionizing and UV radiation.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506585
NCBI GeneID	6118
NCBI Accession #	NP_002937.1
NCBI GenBank Nucleotide #	NM_002946.3
UniProt Accession #	P15927; Q52II0; Q5TEI9; Q5TEJ5
Molecular Weight	29,247 Da
NCBI Official Full Name	replication protein A 32 kDa subunit
NCBI Official Synonym Full Names	replication protein A2, 32kDa
NCBI Official Symbol	RPA2
NCBI Official Synonym Symbols	REPA2; RPA32
NCBI Protein Information	replication protein A 32 kDa subunit; RP-A p32; RP-A p34; RF-A protein 2; replication factor A protein 2; replication protein A 34 kDa subunit
UniProt Protein Name	Replication protein A 32 kDa subunit
UniProt Gene Name	RPA2
UniProt Synonym Gene Names	REPA2; RPA32; RPA34; RP-A p32; RF-A protein 2; RP-A p34
UniProt Entry Name	RFA2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.