

Mouse Thyroglobulin Monoclonal Antibody | anti-TG antibody

Thyroglobulin (Thyroidal Cell Marker) Recombinant Mouse Monoclonal Antibody [Clone rTGB/4744]

CATALOG # AAA215954	HOST Mouse	REACTIVITY Human, Mouse and Rat	APPLICATIONS Immunohistochemistry, Flow Cytometry, Functional Assay
-------------------------------	----------------------	---	---


PRODUCT SYNONYMS

Thyroglobulin; Monoclonal Antibody; Thyroglobulin (Thyroidal Cell Marker) Recombinant Mouse Monoclonal Antibody [Clone rTGB/4744]; Thyroglobulin (Thyroidal Cell Marker); AITD3; hTG; TDH3; Tg; Tgn; anti-TG antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Mouse and Rat
Clonality	Monoclonal
Isotype	Mouse/IgG2a, kappa
Clone Number	rTGB/4744
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Concentration	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml (varies by lot)
Applicable Applications for anti-TG antibody	IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry)
Antibody Type	Recombinant Antibody
Immunogen	Human thyroid follicular cells
Cellular Localization	Cytoplasmic and secreted
Positive Control	Thyroid
Hu-chromosome Location	8q24.22
Unigene	654591
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C Antibody without azide: store at -20 to -80 degree C Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human thyroid carcinoma stained with Thyroglobulin
Mouse Recombinant Monoclonal Antibody (rTGB/4744).

ADDITIONAL INFORMATION

Related Product Information for anti-TG antibody	rTGB/4744 reacts with a partially defined epitope of human thyroglobulin. This epitope is different from the epitope recognized by MAb 6E1. Thyroglobulin is a 660kDa dimeric pre-protein with multiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers undergo conformational maturation in the endoplasmic reticulum. The vast majority of follicular carcinomas of the thyroid will give positive immunoreactivity for anti-thyroglobulin even though sometimes only focally. Poorly differentiated carcinomas of the thyroid are frequently anti-thyroglobulin negative. Adenocarcinomas of other-than-thyroid origin do not react with this antibody. This antibody is useful in identification of thyroid carcinoma of the papillary and follicular types. Presence of thyroglobulin in metastatic lesions establishes the thyroid origin of tumor. Anti-thyroglobulin, combined with anti-calcitonin, can identify medullary carcinomas of the thyroid. Furthermore, anti-thyroglobulin, combined with anti-TTF1, can be a reliable marker to differentiate between primary thyroid and lung neoplasms.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	55770862
NCBI GeneID	7038
NCBI Accession #	NP_003226.4
NCBI GenBank Nucleotide #	NM_003235.4
UniProt Accession #	P01266; O15274; O43899; Q15593; Q15948; Q9NYR1; Q9NYR2; Q9UMZ0; Q9UNY3
Molecular Weight	304,790 Da
NCBI Official Full Name	thyroglobulin
NCBI Official Synonym Full Names	thyroglobulin
NCBI Official Symbol	TG
NCBI Official Synonym Symbols	TGN; AITD3
NCBI Protein Information	thyroglobulin
UniProt Protein Name	Thyroglobulin
UniProt Gene Name	TG
UniProt Synonym Gene Names	Tg
UniProt Entry Name	THYG_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/thyroglobulin-antibody-clone-rtgb-4744-immunohistochemistry-flow-cytometry-elisa/215954>