

Mouse anti-Human TIM3/HAVCR2/CD366 Monoclonal Antibody | anti-HAVCR2 antibody

TIM3/HAVCR2/CD366 (Effector T-Cell Marker) Mouse Monoclonal Antibody [Clone TIM3/4030]

CATALOG # AAA216010	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

TIM3/HAVCR2/CD366; Monoclonal Antibody; TIM3/HAVCR2/CD366 (Effector T-Cell Marker) Mouse Monoclonal Antibody [Clone TIM3/4030]; TIM3/HAVCR2/CD366 (Effector T-Cell Marker); CD366; HAVR2; Hepatitis A virus cellular receptor 2 (HAVCR2); Kidney injury molecule 3 (KIM3); T-cell immunoglobulin and mucin domain-containing protein 3; T-cell immunoglobulin mucin receptor 3; T-cell membrane protein 3; TIM3; TIMD3; anti-HAVCR2 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse/IgG2b, kappa
Clone Number	TIM3/4030
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Concentration	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml (varies by lot)
Applicable Applications for anti-HAVCR2 antibody	IHC (Immunohistochemistry)
Immunogen	Recombinant fragment of human TIM3 protein (around aa 22-202) (exact sequence is proprietary)
Cellular Localization	Cell Surface
Positive Control	HepG2 or Daudi cells. Human tonsil, lymph node or spleen.
Hu-chromosome Location	5q33.3
Unigene	710500
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C Antibody without azide: store at -20 to -80 degree C Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)

Formalin-fixed, paraffin-embedded human lymph node stained with TIM3 Mouse Monoclonal Antibody (TIM3/4030).

ADDITIONAL INFORMATION

Related Product Information for anti-HAVCR2 antibody	TIMs are type I transmembrane glycoproteins with one Ig-like V-type domain and a Ser/Thr-rich mucin stalk. TIM-3 is expressed on the surface of effector T cel tonsil, lymph node or spleen. Is (CD4+Th1 and CD8+Tc1) but not on helper T cells (CD4+Th2 and CD8+Tc2). In chronic inflammation, autoimmune disorders, and some cancers, TIM-3 is upregulated on several other hematopoietic cell types. The Ig domain of TIM-3 interacts with a ligand on resting but not activated Th1 and Th2 cells. The glycosylated Ig domain of TIM-3 binds cell-associated galectin-9. This induces TIM-3 Tyr phosphorylation and pro-apoptotic signaling. TIM-3 functions as a negative regulator of Th1 cell activity. Its blockade results in increased IFN-gamma production, Th1 cell proliferation and cytotoxicity, regulatory T cell development, and increases in macrophage and neutrophil infiltration into sites of inflammation.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	311033536
NCBI GeneID	84868
UniProt Accession #	Q8TDQ0; Q8WW60; Q96K94; B2RAY2
Molecular Weight	301
NCBI Official Full Name	Hepatitis A virus cellular receptor 2
NCBI Official Synonym Full Names	hepatitis A virus cellular receptor 2
NCBI Official Symbol	HAVCR2
NCBI Official Synonym Symbols	TIM3; KIM-3; TIMD3; Tim-3; TIMD-3; HAVcr-2
NCBI Protein Information	hepatitis A virus cellular receptor 2; kidney injury molecule-3; T-cell membrane protein 3; T cell immunoglobulin mucin 3; T cell immunoglobulin mucin-3; T-cell immunoglobulin mucin receptor 3; T-cell immunoglobulin mucin family member 3; T-cell immunoglo
UniProt Protein Name	Hepatitis A virus cellular receptor 2
UniProt Gene Name	HAVCR2
UniProt Synonym Gene Names	TIM3; TIMD3; HAVcr-2; TIMD-3; TIM-3
UniProt Entry Name	HAVR2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.