

Mouse anti-Human CD5 Monoclonal Antibody | anti-CD5 antibody

CD5 (Mantle Cell Lymphoma Marker) Recombinant Mouse Monoclonal Antibody [Clone rC5/6462]

CATALOG # AAA216035	HOST Mouse	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	----------------------	----------------------------	---


PRODUCT SYNONYMS

CD5; Monoclonal Antibody; CD5 (Mantle Cell Lymphoma Marker) Recombinant Mouse Monoclonal Antibody [Clone rC5/6462]; CD5 (Mantle Cell Lymphoma Marker); CD5 antigen (p56 62); LEU1; Ly12; LyA; Lymphocyte antigen T1/Leu-1; T-cell surface glycoprotein CD5; anti-CD5 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse/IgG2a, kappa
Clone Number	rC5/6462
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Concentration	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml (varies by lot)
Applicable Applications for anti-CD5 antibody	IHC (Immunohistochemistry)
Antibody Type	Recombinant Antibody
Immunogen	Recombinant full-length protein corresponding to human CD5
Cellular Localization	Cell Surface
Positive Control	293T, Ramos, or MOLT-4 cells. Human tonsil or lymph node.
Hu-chromosome Location	11q12.2
Unigene	58685
Preparation and Storage	Antibody with azide: store at 2 to 8 degree C Antibody without azide: store at -20 to -80 degree C Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)

Formalin-fixed, paraffin-embedded human tonsil stained with CD5 Recombinant Mouse Monoclonal Antibody (rC5/6462).

ADDITIONAL INFORMATION

Related Product Information for anti-CD5 antibody	The CD5 antigen is a transmembrane glycoprotein expressed on the surface of practically all mature human T-cells (about 10% of CD4+ T-cells being CD5 negative). In immature (CD34+) T-cells, CD5 is weakly expressed, the intensity of expression increasing with maturation. CD5 is also expressed in a small subset of normal human B-cells (20% of B-cells in the peripheral blood, scattered cells in the lymph node mantle zone). The CD5+ cells are probably involved in B-T interaction and their ligand is CD72 which is expressed on all B cells. It appears that CD5+ B-cells on activation primarily produce IgM. They also produce more autoantibodies than normal CD5 negative B-cells. Thus, the CD5+ B-cell population is expanded in rheumatoid arthritis and systemic lupus erythematosus. The CD5 antibody reacts with CD5 expressed on B and T cells, and may be a useful aid for the classification of B and T-cell malignancies. This includes B-cell chronic lymphoid leukemia (B-CLL), B-cell small lymphocytic lymphoma (B-SLL), mantle cell lymphoma (MCL) and T-cell lymphoma and leukemia.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	166197668
NCBI GeneID	921
NCBI Accession #	NP_055022.2
NCBI GenBank Nucleotide #	NM_014207.3
UniProt Accession #	P06127; A0N0P4; A8K9I3
Molecular Weight	54,578 Da
NCBI Official Full Name	T-cell surface glycoprotein CD5
NCBI Official Synonym Full Names	CD5 molecule
NCBI Official Symbol	CD5
NCBI Official Synonym Symbols	T1; LEU1
NCBI Protein Information	T-cell surface glycoprotein CD5; CD5 antigen (p56-62); lymphocyte antigen T1/Leu-1
UniProt Protein Name	T-cell surface glycoprotein CD5
UniProt Gene Name	CD5
UniProt Synonym Gene Names	LEU1
UniProt Entry Name	CD5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd5-antibody-clone-rc5-6462-immunohistochemistry-elisa/216035>