

Human Inducible T-Cell Co Stimulator Ligand (ICOSLG) RTU ELISA Kit | ICOSLG rtu elisa kit

Human Inducible T-Cell Co Stimulator Ligand (ICOSLG) ELISA Kit

CATALOG #
AAA216120

REACTIVITY
Human


PRODUCT SYNONYMS

Inducible T-Cell Co Stimulator Ligand (ICOSLG); Human Inducible T-Cell Co Stimulator Ligand (ICOSLG) ELISA Kit; CD275; B7-H2; B7H2; B7RP-1; B7RP1; GL50; ICOS-L; ICOSL; LICOS; B7-like protein GL50; B7-related protein 1; ICOSLG rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Sequence Length	309
Assay Type	Sandwich
Samples	Tissue Homogenates, Cell Lysates or Other Biological Fluids.
Detection Range	0.156-10ng/mL
Sensitivity	0.057ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ICOSLG rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to ICOSLG. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to ICOSLG. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ICOSLG, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of ICOSLG in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	545688560
NCBI GeneID	23308
NCBI Accession #	NP_001269979.1
NCBI GenBank Nucleotide #	NM_001283050.1
UniProt Accession #	O75144; Q9HD18; Q9NRQ1; A8MUZ1
Molecular Weight	20,120 Da
NCBI Official Full Name	ICOS ligand isoform b
NCBI Official Synonym Full Names	inducible T cell costimulator ligand
NCBI Official Symbol	ICOSLG
NCBI Official Synonym Symbols	B7H2; GL50; B7-H2; B7RP1; CD275; ICOSL; LICOS; B7RP-1; ICOS-L
NCBI Protein Information	ICOS ligand
UniProt Protein Name	ICOS ligand
UniProt Gene Name	ICOSLG
UniProt Synonym Gene Names	B7H2; B7RP1; ICOSL; KIAA0653; B7-H2; B7RP-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-rtu-elisa-kits/inducible-t-cell-co-stimulator-ligand-icoslg/216120>