

Human Interleukin 2 Receptor Gamma (IL2Rg) RTU ELISA Kit | IL2Rg rtu elisa kit

Human Interleukin 2 Receptor Gamma (IL2Rg) ELISA Kit

CATALOG #
AAA216403

REACTIVITY
Human


PRODUCT SYNONYMS

Interleukin 2 Receptor Gamma (IL2Rg); Human Interleukin 2 Receptor Gamma (IL2Rg) ELISA Kit; CD132; IL2R-g; IL2-RG; IMD4; p64; SCIDX; SCIDX1; Cytokine Receptor Common Gamma Chain; Severe Combined Immunodeficiency; IL2Rg rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Sequence Length	369
Assay Type	Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates or Other Biological Fluids.
Detection Range	0.156-10ng/mL
Sensitivity	0.067ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for IL2Rg rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to IL2Rg. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to IL2Rg. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain IL2Rg, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of IL2Rg in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557882
NCBI GeneID	3561
NCBI Accession #	NP_000197.1
NCBI GenBank Nucleotide #	NM_000206.2
UniProt Accession #	P31785; Q5FC12
Molecular Weight	20,088 Da
NCBI Official Full Name	cytokine receptor common subunit gamma
NCBI Official Synonym Full Names	interleukin 2 receptor subunit gamma
NCBI Official Symbol	IL2RG
NCBI Official Synonym Symbols	P64; CIDX; IMD4; CD132; SCIDX; IL-2RG; SCIDX1
NCBI Protein Information	cytokine receptor common subunit gamma
UniProt Protein Name	Cytokine receptor common subunit gamma
UniProt Gene Name	IL2RG
UniProt Synonym Gene Names	IL-2 receptor subunit gamma; IL-2R subunit gamma; IL-2RG

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-rtu-elisa-kits/interleukin-2-receptor-gamma-il2rg/216403>