

Mouse Melanoma Cell Adhesion Molecule (MCAM) RTU ELISA Kit | MCAM rtu elisa kit

Mouse Melanoma Cell Adhesion Molecule (MCAM) ELISA Kit

CATALOG #
AAA216956

REACTIVITY
Mouse


PRODUCT SYNONYMS

Melanoma Cell Adhesion Molecule (MCAM); Mouse Melanoma Cell Adhesion Molecule (MCAM) ELISA Kit; CD146; MUC18; Cell surface glycoprotein P1H12; Melanoma-associated antigen A32; S-endo 1 endothelial-associated antigen; MCAM rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Sequence Length	648
Assay Type	Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates or Other Biological Fluids.
Detection Range	0.625-40ng/mL
Sensitivity	0.234ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MCAM rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to MCAM. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to MCAM. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain MCAM, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of MCAM in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	160333901
NCBI GeneID	84004
NCBI Accession #	NP_075548.2
NCBI GenBank Nucleotide #	NM_023061.2
UniProt Accession #	Q8R2Y2; Q9EPF1; Q9ESS7; Q9JHQ2; Q9JHQ3
Molecular Weight	66,901 Da
NCBI Official Full Name	cell surface glycoprotein MUC18 isoform 1
NCBI Official Synonym Full Names	melanoma cell adhesion molecule
NCBI Official Symbol	Mcam
NCBI Official Synonym Symbols	CD146; CD149; Muc18; s-endo; AV025631; 1-gicerin; s-gicerin
NCBI Protein Information	cell surface glycoprotein MUC18
UniProt Protein Name	Cell surface glycoprotein MUC18
UniProt Gene Name	Mcam
UniProt Synonym Gene Names	Muc18

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-rtu-elisa-kits/melanoma-cell-adhesion-molecule-mcam/216956>