

Mouse Monokine Induced By Interferon Gamma (Mlg) RTU ELISA Kit | Mlg rtu elisa kit

Mouse Monokine Induced By Interferon Gamma (Mlg) ELISA Kit

CATALOG #
AAA217039

REACTIVITY
Mouse


PRODUCT SYNONYMS

Monokine Induced By Interferon Gamma (Mlg); Mouse Monokine Induced By Interferon Gamma (Mlg) ELISA Kit; CXCL9; CMK; Humig; MI-G; SCYB9; Crg-10; Chemokine (C-X-C motif) ligand 9; Small-inducible cytokine B9; Gamma-interferon-induced monokine; Mlg rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Sequence Length	126
Assay Type	Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates or Other Biological Fluids.
Detection Range	46.8-3000pg/mL
Sensitivity	18.35pg/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for Mlg rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to Mlg. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to Mlg. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Mlg, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of Mlg in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	110626107
NCBI GeneID	17329
NCBI Accession #	NP_032625.2
NCBI GenBank Nucleotide #	NM_008599.4
UniProt Accession #	P18340; Q8C9J0; Q99J60
Molecular Weight	14,444 Da
NCBI Official Full Name	C-X-C motif chemokine 9
NCBI Official Synonym Full Names	chemokine (C-X-C motif) ligand 9
NCBI Official Symbol	Cxcl9
NCBI Official Synonym Symbols	CMK; Mig; MuMIG; Scyb9; crg-10; BB139920
NCBI Protein Information	C-X-C motif chemokine 9
UniProt Protein Name	C-X-C motif chemokine 9
UniProt Gene Name	Cxcl9
UniProt Synonym Gene Names	Mig; Scyb9; MIG; MuMIG

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-rtu-elisa-kits/monokine-induced-by-interferon-gamma-mig/217039>