

Human N-Methylpurine DNA Glycosylase (MPG) RTU ELISA Kit | MPG rtu elisa kit

Human N-Methylpurine DNA Glycosylase (MPG) ELISA Kit

CATALOG #
AAA217149

REACTIVITY
Human


PRODUCT SYNONYMS

N-Methylpurine DNA Glycosylase (MPG); Human N-Methylpurine DNA Glycosylase (MPG) ELISA Kit; AAG; APNG; ADPG; CRA36.1; MDG; Mid1; PIG11; PIG16; Anpg; DNA-3-Methyladenine Glycosylase; Alkyladenine DNA Glycosylase; 3-alkyladenine DNA glycosylase; MPG rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Sequence Length	293
Assay Type	Sandwich
Samples	Tissue Homogenates or Other Biological Fluids.
Detection Range	0.156-10ng/mL
Sensitivity	0.055ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MPG rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to MPG. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to MPG. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain MPG, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of MPG in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	62632765
NCBI GeneID	4350

NCBI Accession #	NP_001015052.1
NCBI GenBank Nucleotide #	NM_001015052.2
UniProt Accession #	P29372 ; Q13770 ; Q15275 ; Q15961 ; Q5J9I4 ; Q96BZ6 ; Q96S33 ; Q9NNX5 ; G5E9E2
Molecular Weight	30,824 Da
NCBI Official Full Name	DNA-3-methyladenine glycosylase isoform b
NCBI Official Synonym Full Names	N-methylpurine DNA glycosylase
NCBI Official Symbol	MPG
NCBI Official Synonym Symbols	AAG; MDG; ADPG; APNG; Mid1; anpg; PIG11; PIG16; CRA36.1
NCBI Protein Information	DNA-3-methyladenine glycosylase
UniProt Protein Name	DNA-3-methyladenine glycosylase
UniProt Gene Name	MPG
UniProt Synonym Gene Names	AAG; ANPG; MID1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/human-rtu-elisa-kits/n-methylpurine-dna-glycosylase-mpg/217149>