

Human Mannose Receptor C Type 2 (MRC2) RTU ELISA Kit | MRC2 rtu elisa kit

Human Mannose Receptor C Type 2 (MRC2) ELISA Kit

CATALOG #
AAA217169

REACTIVITY
Human


PRODUCT SYNONYMS

Mannose Receptor C Type 2 (MRC2); Human Mannose Receptor C Type 2 (MRC2) ELISA Kit; CD280; ENDO180; CLEC13E; UPARAP; C-type lectin domain family 13 member E; Endocytic receptor 180; Macrophage mannose receptor 2; Urokinase receptor-associated protein; MRC2 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Sequence Length	1479
Assay Type	Sandwich
Samples	Serum, Plasma, Tissue Homogenates or Other Biological Fluids.
Detection Range	0.312-20ng/mL
Sensitivity	0.117ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MRC2 rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to MRC2. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to MRC2. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain MRC2, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of MRC2 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	110624774
NCBI GeneID	9902
NCBI Accession #	NP_006030.2
NCBI GenBank Nucleotide #	NM_006039.4
UniProt Accession #	Q9UBG0; Q7LGE7; Q9Y5P9; A6H8K4; D3DU08
Molecular Weight	166,674 Da
NCBI Official Full Name	C-type mannose receptor 2
NCBI Official Synonym Full Names	mannose receptor C type 2
NCBI Official Symbol	MRC2
NCBI Official Synonym Symbols	CD280; UPARAP; CLEC13E; ENDO180
NCBI Protein Information	C-type mannose receptor 2
UniProt Protein Name	C-type mannose receptor 2
UniProt Gene Name	MRC2
UniProt Synonym Gene Names	CLEC13E; ENDO180; KIAA0709; UPARAP; UPAR-associated protein; Urokinase receptor-associated protein

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/human-rtu-elisa-kits/mannose-receptor-c-type-2-mrc2/217169>