

Rat V-Myc Myelocytomatosis Viral Oncogene Homolog (MYC) RTU ELISA Kit | MYC rtu elisa kit

Rat V-Myc Myelocytomatosis Viral Oncogene Homolog (MYC) ELISA Kit

CATALOG #
AAA217248

REACTIVITY
Rat


PRODUCT SYNONYMS

V-Myc Myelocytomatosis Viral Oncogene Homolog (MYC); Rat V-Myc Myelocytomatosis Viral Oncogene Homolog (MYC) ELISA Kit; C-Myc; BHLHE39; Myc Proto-Oncogene Protein; Class E basic helix-loop-helix protein 39; Transcription factor p64; Proto-oncogene c-Myc; MYC rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Sequence Length	453
Assay Type	Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates or Other Biological Fluids.
Detection Range	0.156-10ng/mL
Sensitivity	0.062ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for MYC rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to MYC. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to MYC. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain MYC, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of MYC in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	71834866
NCBI GeneID	24577
NCBI Accession #	NP_036735.2
NCBI GenBank Nucleotide #	NM_012603.2
UniProt Accession #	P09416; Q6B500
Molecular Weight	48,898 Da
NCBI Official Full Name	myc proto-oncogene protein
NCBI Official Synonym Full Names	MYC proto-oncogene, bHLH transcription factor
NCBI Official Symbol	Myc
NCBI Official Synonym Symbols	mMyc; c-myc; RNCMYC
NCBI Protein Information	myc proto-oncogene protein
UniProt Protein Name	Myc proto-oncogene protein
UniProt Gene Name	Myc

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-rtu-elisa-kits/v-myc-myelocytomatosis-viral-oncogene-homolog-myc/217248>