

Mouse Norepinephrine Transporter (NET) RTU ELISA Kit | NET rtu elisa kit

Mouse Norepinephrine Transporter (NET) ELISA Kit

CATALOG #
AAA217343

REACTIVITY
Mouse


PRODUCT SYNONYMS

Norepinephrine Transporter (NET); Mouse Norepinephrine Transporter (NET) ELISA Kit; SLC6A2; NAT1; NET; NET1; Sodium-dependent noradrenaline transporter; Noradrenaline Transporter; Solute Carrier Family 6 member 2; Neurotransmitter transporter; Noradrenalin; NET rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Sequence Length	617
Assay Type	Sandwich
Samples	Tissue Homogenates, Cell Lysates or Other Biological Fluids.
Detection Range	0.156-10ng/mL
Sensitivity	0.062ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for NET rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to NET. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to NET. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain NET, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of NET in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	165377298
NCBI GeneID	20538
NCBI Accession #	NP_033235.3
NCBI GenBank Nucleotide #	NM_009209.3
UniProt Accession #	O55192; Q6QU62
Molecular Weight	69,255 Da
NCBI Official Full Name	sodium-dependent noradrenaline transporter
NCBI Official Synonym Full Names	solute carrier family 6 (neurotransmitter transporter, noradrenalin), member 2
NCBI Official Symbol	Slc6a2
NCBI Official Synonym Symbols	NET; NE-T; Slc6a5
NCBI Protein Information	sodium-dependent noradrenaline transporter
UniProt Protein Name	Sodium-dependent noradrenaline transporter
UniProt Gene Name	Slc6a2
UniProt Synonym Gene Names	NET

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-rtu-elisa-kits/norepinephrine-transporter-net/217343>