

Mouse Poly ADP Ribose Polymerase (PARP) RTU ELISA Kit | PARP rtu elisa kit

Mouse Poly ADP Ribose Polymerase (PARP) ELISA Kit

CATALOG #
AAA217644

REACTIVITY
Mouse


PRODUCT SYNONYMS

Poly ADP Ribose Polymerase (PARP); Mouse Poly ADP Ribose Polymerase (PARP) ELISA Kit; PARP1; ADPRT; ADPRT1; PPOL; pADPRT-1; ADP-ribosyltransferase diphtheria toxin-like 1; NAD(+) ADP-ribosyltransferase 1; PARP rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of PARP. No significant cross-reactivity or interference between PARP and analogues was observed.
Sequence Length	1013
Samples	Serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	31.2-2000pg/mL
Sensitivity	< 13.26pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level PARP were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level PARP were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PARP rtu elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of PARP in mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to PARP. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to PARP. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain PARP, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of PARP in the samples is then determined by comparing the O.D. of the samples to the standard curve.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	130782
NCBI GeneID	11545
UniProt Accession #	P11103; Q9JLX4; Q9QVQ3
Molecular Weight	55,237 Da
NCBI Official Full Name	Poly
NCBI Official Synonym Full Names	poly (ADP-ribose) polymerase family, member 1
NCBI Official Symbol	Parp1
NCBI Official Synonym Symbols	PARP; PPOL; ARTD1; Adprp; Adprt1; C80510; msPARP; parp-1; sPARP-1; AI893648; 5830444G22Rik
NCBI Protein Information	poly [ADP-ribose] polymerase 1
UniProt Protein Name	Poly [ADP-ribose] polymerase 1
UniProt Gene Name	Parp1
UniProt Synonym Gene Names	Adprp; Adprt; Adprt1; PARP-1; ARTD1; ADPRT 1; msPARP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.