

Mouse Peroxisome Proliferator Activated Receptor Alpha (PPARα) RTU ELISA Kit | PPARα rtu elisa kit

Mouse Peroxisome Proliferator Activated Receptor Alpha (PPARα) ELISA Kit

CATALOG #
AAA217934

REACTIVITY
Mouse


PRODUCT SYNONYMS

Peroxisome Proliferator Activated Receptor Alpha (PPARα); Mouse Peroxisome Proliferator Activated Receptor Alpha (PPARα) ELISA Kit; NR1C1; PPAR-A; HPPAR; Nuclear Receptor Subfamily 1 Group C Member 1; PPARα rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Sequence Length	468
Assay Type	Sandwich
Samples	Serum, Plasma, Tissue Homogenates, Cell Lysates, Cell Culture Supernates or Other Biological Fluids.
Detection Range	0.156-10ng/mL
Sensitivity	0.051ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for PPARα rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to PPARα. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to PPARα. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain PPARα, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of PPARα in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	164663880
NCBI GeneID	19013
NCBI Accession #	NP_001106889.1
NCBI GenBank Nucleotide #	NM_001113418.1
UniProt Accession #	P23204
Molecular Weight	52,347 Da
NCBI Official Full Name	peroxisome proliferator-activated receptor alpha
NCBI Official Synonym Full Names	peroxisome proliferator activated receptor alpha
NCBI Official Symbol	Ppara
NCBI Official Synonym Symbols	Ppar; Nr1c1; AW742785; PPARalpha; PPAR-alpha; 4933429D07Rik
NCBI Protein Information	peroxisome proliferator-activated receptor alpha
UniProt Protein Name	Peroxisome proliferator-activated receptor alpha
UniProt Gene Name	Ppara
UniProt Synonym Gene Names	Nr1c1; Ppar; PPAR-alpha

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-rtu-elisa-kits/peroxisome-proliferator-activated-receptor-alpha-ppara/217934>