

Rat Sirtuin 3 (SIRT3) RTU ELISA Kit | SIRT3 rtu elisa kit

Rat Sirtuin 3 (SIRT3) ELISA Kit

CATALOG #
AAA218433

REACTIVITY
Rat


PRODUCT SYNONYMS

Sirtuin 3 (SIRT3); Rat Sirtuin 3 (SIRT3) ELISA Kit; SIR2L3; Silent Mating Type Information Regulation 2 Homolog 3; NAD-dependent protein deacetylase sirtuin-3; mitochondrial; SIR2-like protein 3; Regulatory protein SIR2 homolog 3; SIRT3 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Rat
Sequence Length	257
Assay Type	Sandwich
Samples	Tissue Homogenates, Cell Lysates or Other Biological Fluids.
Detection Range	0.312-20ng/mL
Sensitivity	0.088ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for SIRT3 rtu elisa kit

Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to SIRT3. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to SIRT3. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain SIRT3, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of SIRT3 in the samples is then determined by comparing the O.D. of the samples to the standard curve.

NCBI AND UNIPROT INFORMATION

NCBI GI #	157821285
NCBI GeneID	293615
NCBI Accession #	NP_001099783.1
NCBI GenBank Nucleotide #	NM_001106313.2
UniProt Accession #	B2RZ31
Molecular Weight	28,497 Da
NCBI Official Full Name	NAD-dependent protein deacetylase sirtuin-3, mitochondrial
NCBI Official Synonym Full Names	sirtuin 3
NCBI Official Symbol	Sirt3
NCBI Protein Information	NAD-dependent protein deacetylase sirtuin-3, mitochondrial
UniProt Protein Name	NAD-dependent protein deacetylase
UniProt Gene Name	Sirt3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rat-rtu-elisa-kits/sirtuin-3-sirt3/218433>