

Mouse Tumor Necrosis Factor Receptor Superfamily, Member 17 (TNFRSF17) RTU ELISA Kit | TNFRSF17 rtu elisa kit

Mouse Tumor Necrosis Factor Receptor Superfamily, Member 17 (TNFRSF17) ELISA Kit

CATALOG # AAA218889	REACTIVITY Mouse
-------------------------------	----------------------------


PRODUCT SYNONYMS

Tumor Necrosis Factor Receptor Superfamily; Member 17 (TNFRSF17); Mouse Tumor Necrosis Factor Receptor Superfamily; Member 17 (TNFRSF17) ELISA Kit; CD269; BCM; BCMA; B-cell maturation protein; TNFRSF17 rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Sequence Length	185
Assay Type	Sandwich
Samples	Serum, Plasma or Other Biological Fluids.
Detection Range	0.312-20ng/mL
Sensitivity	0.101ng/mL
RTU ELISA Kit's Advantage	The Ready-To-Use (RTU) Kits require fewer steps in setup and take less time to run. Detection Solutions are pre-diluted and total incubation time is shorter than many other ELISA Kits.
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TNFRSF17 rtu elisa kit	Test Principle: The microtiter plate provided in this kit has been pre-coated with an antibody specific to TNFRSF17. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to TNFRSF17. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain TNFRSF17, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm $\pm$ 10nm. The concentration of TNFRSF17 in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	6755823
-----------	---------

NCBI GeneID	21935
NCBI Accession #	NP_035738.1
NCBI GenBank Nucleotide #	NM_011608.1
UniProt Accession #	O88472
Molecular Weight	19,932 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 17
NCBI Official Synonym Full Names	tumor necrosis factor receptor superfamily, member 17
NCBI Official Symbol	Tnfrsf17
NCBI Official Synonym Symbols	BCM; BCMA; Tnfrsf13; Tnfrsf13a
NCBI Protein Information	tumor necrosis factor receptor superfamily member 17
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 17
UniProt Gene Name	Tnfrsf17
UniProt Synonym Gene Names	Bcm; Bcma

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/mouse-rtu-elisa-kits/tumor-necrosis-factor-receptor-superfamily-member-17-tnfrsf17/218889>