

Human Vinculin (VCL) RTU ELISA Kit | VCL rtu elisa kit

Human Vinculin (VCL) ELISA Kit

CATALOG #
AAA219147

REACTIVITY
Human


PRODUCT SYNONYMS

Vinculin (VCL); Human Vinculin (VCL) ELISA Kit; MVCL; Metavinculin; VCL rtu elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of VCL. No significant cross-reactivity or interference between VCL and analogues was observed.
Sequence Length	1066
Samples	Serum, plasma, tissue homogenates or other biological fluids
Assay Type	Quantitative Sandwich
Detection Range	78.1-5000pg/mL
Sensitivity	< 32pg/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level VCL were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level VCL were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	For unopened kits: All the reagents should be kept at 4 degree C upon receipt. For opened kits: Once the kit is opened, the remaining reagents still need to be stored according to the above storage conditions. In addition, return the unused wells to the foil pouch containing the desiccant pack and reseal along entire edge of zip-seal.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for VCL rtu elisa kit	Intended Uses: The kit is a sandwich enzyme immunoassay for the in vitro quantitative measurement of VCL in human serum, plasma, tissue homogenates or other biological fluids. Principle of the Assay: The microtiter plate provided in this kit has been pre-coated with an antibody specific to VCL. Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody preparation specific to VCL. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain VCL, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of VCL in the samples is then determined by comparing the O.D. of the samples to the standard curve.
---	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507877
NCBI GeneID	7414
NCBI Accession #	NP_003364.1
NCBI GenBank Nucleotide #	NM_003373.3
UniProt Accession #	P18206; Q16450; Q5SWX2; Q7Z3B8; Q8IXU7
Molecular Weight	24,904 Da
NCBI Official Full Name	vinculin isoform VCL
NCBI Official Synonym Full Names	vinculin
NCBI Official Symbol	VCL
NCBI Official Synonym Symbols	MV; MVCL; CMD1W; CMH15; HEL114
NCBI Protein Information	vinculin
UniProt Protein Name	Vinculin
UniProt Gene Name	VCL
UniProt Synonym Gene Names	MV

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.