

Rabbit anti-Human Cytochrome P450 27A1 Polyclonal Antibody | anti-CYP27A1 antibody

Rabbit anti-Cytochrome P450 27A1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220303	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Cytochrome P450 27A1; Polyclonal Antibody; Rabbit anti-Cytochrome P450 27A1 Antibody; CYP27; Sterol 26-hydroxylase; mitochondrial; 5-beta-cholestane-3-alpha; 7-alpha; 12-alpha-triol 27-hydroxylase; Cytochrome P-450C27/25; Cytochrome P450 27; Sterol 27-hydroxylase; Vitamin D(3) 25-hydroxylase; anti-CYP27A1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Clone Number	EH12.2H7
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CYP27A1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Cytochrome P450 27A1. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Cytochrome P450 27A1 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Cytochrome P450 27A1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of Cytochrome P450 27A1 expression in HepG2 (A), Caco2 (B), A549 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CYP27A1 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503211
NCBI GeneID	1593
NCBI Accession #	NP_000775.1
NCBI GenBank Nucleotide #	NM_000784.3
UniProt Accession #	Q02318; Q6LDB4; Q86YQ6; A8K303
Molecular Weight	60,235 Da
NCBI Official Full Name	sterol 26-hydroxylase, mitochondrial
NCBI Official Synonym Full Names	cytochrome P450, family 27, subfamily A, polypeptide 1
NCBI Official Symbol	CYP27A1
NCBI Official Synonym Symbols	CTX; CP27; CYP27
NCBI Protein Information	sterol 26-hydroxylase, mitochondrial; cytochrome P450 27; sterol 27-hydroxylase; cytochrome P-450C27/25; vitamin D(3) 25-hydroxylase; cholestanetriol 26-monooxygenase; 5-beta-cholestane-3-alpha,7-alpha,12-alpha-triol 27-hydroxylase; 5-beta-cholestane-3-al
UniProt Protein Name	Sterol 26-hydroxylase, mitochondrial
UniProt Gene Name	CYP27A1
UniProt Synonym Gene Names	CYP27
UniProt Entry Name	CP27A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.