

Rabbit anti-Human MCM2 Polyclonal Antibody | anti-MCM2 antibody

Rabbit anti-MCM2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220327	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MCM2; Polyclonal Antibody; Rabbit anti-MCM2 Antibody; BM28; CCNL1; CDCL1; KIAA0030; DNA replication licensing factor MCM2; Minichromosome maintenance protein 2 homolog; Nuclear protein BM28; anti-MCM2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-MCM2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human MCM2. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MCM2 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of MCM2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MCM2 expression in HeLa (A), Jurkat (B), HeLa (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-MCM2 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	33356547
NCBI GeneID	4171
NCBI Accession #	NP_004517.2
NCBI GenBank Nucleotide #	NM_004526.3
UniProt Accession #	P49736; Q14577; Q15023; Q8N2V1; Q969W7; Q96AE1; Q9BRM7
Molecular Weight	101,896 Da
NCBI Official Full Name	DNA replication licensing factor MCM2
NCBI Official Synonym Full Names	minichromosome maintenance complex component 2
NCBI Official Symbol	MCM2
NCBI Official Synonym Symbols	BM28; CCNL1; CDCL1; cdc19; D3S3194; MITOTIN
NCBI Protein Information	DNA replication licensing factor MCM2; cyclin-like 1; nuclear protein BM28; cell division cycle-like 1; minichromosome maintenance protein 2 homolog; minichromosome maintenance deficient 2 (mitotin); MCM2 minichromosome maintenance deficient 2, mitotin
UniProt Protein Name	DNA replication licensing factor MCM2
UniProt Gene Name	MCM2
UniProt Synonym Gene Names	BM28; CCNL1; CDCL1; KIAA0030
UniProt Entry Name	MCM2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/mcm2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220327>