

Rabbit anti-Human PKR Polyclonal Antibody | anti-PKR antibody

Rabbit anti-PKR Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220339	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PKR; Polyclonal Antibody; Rabbit anti-PKR Antibody; PRKR; Interferon-induced; double-stranded RNA-activated protein kinase; Eukaryotic translation initiation factor 2-alpha kinase 2; eIF-2A protein kinase 2; Interferon-inducible RNA-dependent protein kinase; P1/eIF-2A protein kinase; Protein kinase RNA-activated; Tyrosine-protein kinase EIF2AK2; p68 kinase; anti-PKR antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-PKR antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PKR. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of PKR staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of PKR staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of PKR expression in HeLa (A), MCF7 (B), Caco2 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-PKR antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	208431827
NCBI GeneID	5610
NCBI Accession #	NP_001129123.1
NCBI GenBank Nucleotide #	NM_001135651.2
UniProt Accession #	P19525; Q52M43; Q7Z6F6; Q9UIR4; A8K3P0; D6W584; E9PC80
Molecular Weight	62,094 Da
NCBI Official Full Name	interferon-induced, double-stranded RNA-activated protein kinase isoform a
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 2-alpha kinase 2
NCBI Official Symbol	EIF2AK2
NCBI Official Synonym Symbols	PKR; PRKR; EIF2AK1
NCBI Protein Information	interferon-induced, double-stranded RNA-activated protein kinase; p68 kinase; eIF-2A protein kinase 2; P1/eIF-2A protein kinase; tyrosine-protein kinase EIF2AK2; interferon-inducible eIF2alpha kinase; double stranded RNA activated protein kinase; protein
UniProt Protein Name	Interferon-induced, double-stranded RNA-activated protein kinase
UniProt Gene Name	EIF2AK2
UniProt Synonym Gene Names	PKR; PRKR; eIF-2A protein kinase 2; PKR
UniProt Entry Name	E2AK2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/pkr-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220339>