

Rabbit anti-Human MGCACGAP Polyclonal Antibody | anti-MGCACGAP antibody

Rabbit anti-MGCACGAP Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220382	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

MGCACGAP; Polyclonal Antibody; Rabbit anti-MGCACGAP Antibody; KIAA1478; Rac GTPase-activating protein 1; Male germ cell RacGap; MgcRacGAP; Protein CYK4 homolog; CYK4; HsCYK-4; anti-MGCACGAP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-MGCACGAP antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human MGCACGAP. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of MGCACGAP staining in K562 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of MGCACGAP staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of MGCRCAGAP expression in Raji (A), K562 (B), A431 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-MGCRCAGAP antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	186910300
NCBI GeneID	29127
NCBI Accession #	NP_001119575.1
NCBI GenBank Nucleotide #	NM_001126103.1
UniProt Accession #	Q9H0H5; Q6PJ26; Q9NWN2; Q9P250; Q9P2W2
Molecular Weight	71,027 Da
NCBI Official Full Name	rac GTPase-activating protein 1
NCBI Official Synonym Full Names	Rac GTPase activating protein 1
NCBI Official Symbol	RACGAP1
NCBI Official Synonym Symbols	CYK4; ID-GAP; HsCYK-4; MgcRacGAP
NCBI Protein Information	rac GTPase-activating protein 1; protein CYK4 homolg; protein CYK4 homolog; male germ cell RacGap; GTPase activating protein
UniProt Protein Name	Rac GTPase-activating protein 1
UniProt Gene Name	RACGAP1
UniProt Synonym Gene Names	KIAA1478; MGCRCAGAP; MgcRacGAP; CYK4; HsCYK-4
UniProt Entry Name	RGAP1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mgcracgap-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220382>