

Rabbit anti-Human COX4-1 Polyclonal Antibody | anti-COX4-1 antibody

Rabbit anti-COX4-1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220452	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

COX4-1; Polyclonal Antibody; Rabbit anti-COX4-1 Antibody; COX4; Cytochrome c oxidase subunit 4 isoform 1; mitochondrial; Cytochrome c oxidase polypeptide IV; Cytochrome c oxidase subunit IV isoform 1; COX IV-1; anti-COX4-1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-COX4-1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human COX4-1. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of COX4-1 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of COX4-1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of COX4-1 expression in HepG2 (A), Jurkat (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-COX4-1 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502981
NCBI GeneID	1327
NCBI Accession #	NP_001852.1
NCBI GenBank Nucleotide #	NM_001861.3
UniProt Accession #	P13073; Q6P666; B2R4J2; D3DUM7
Molecular Weight	19,577 Da
NCBI Official Full Name	cytochrome c oxidase subunit 4 isoform 1, mitochondrial
NCBI Official Synonym Full Names	cytochrome c oxidase subunit IV isoform 1
NCBI Official Symbol	COX4I1
NCBI Official Synonym Symbols	COX4; COXIV; COX4-1
NCBI Protein Information	cytochrome c oxidase subunit 4 isoform 1, mitochondrial; COX IV-1; cytochrome c oxidase polypeptide IV
UniProt Protein Name	Cytochrome c oxidase subunit 4 isoform 1, mitochondrial
UniProt Gene Name	COX4I1
UniProt Synonym Gene Names	COX4; COX IV-1
UniProt Entry Name	COX41_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cox4-1-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220452>