

Rabbit anti-Human RAD17 Polyclonal Antibody | anti-RAD17 antibody

Rabbit anti-RAD17 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220481	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

RAD17; Polyclonal Antibody; Rabbit anti-RAD17 Antibody; R24L; Cell cycle checkpoint protein RAD17; hRad17; RF-C/activator 1 homolog; anti-RAD17 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-RAD17 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human RAD17. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of RAD17 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of RAD17 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of RAD17 expression in human Oral cancer (A), HuvEc (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-RAD17 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506383
NCBI GeneID	5884
NCBI Accession #	NP_002864.1
NCBI GenBank Nucleotide #	NM_002873.1
UniProt Accession #	O75943; O75714; Q7Z3S4; Q9UNK7; Q9UNR7; Q9UNR8; Q9UPF5; A8K8X2; D3DWA5
Molecular Weight	77,055 Da
NCBI Official Full Name	cell cycle checkpoint protein RAD17 isoform 1
NCBI Official Synonym Full Names	RAD17 homolog (S. pombe)
NCBI Official Symbol	RAD17
NCBI Official Synonym Symbols	CCYC; R24L; RAD24; HRAD17; RAD17SP
NCBI Protein Information	cell cycle checkpoint protein RAD17; RAD1 homolog; Rad17-like protein; RF-C activator 1 homolog; RF-C/activator 1 homolog
UniProt Protein Name	Cell cycle checkpoint protein RAD17
UniProt Gene Name	RAD17
UniProt Synonym Gene Names	R24L; hRad17
UniProt Entry Name	RAD17_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/rad17-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220481>