

Rabbit anti-Human CD254 Polyclonal Antibody | anti-CD254 antibody

Rabbit anti-CD254 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220489	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CD254; Polyclonal Antibody; Rabbit anti-CD254 Antibody; OPGL; RANKL; TRANCE; Tumor necrosis factor ligand superfamily member 11; Osteoclast differentiation factor; ODF; Osteoprotegerin ligand; Receptor activator of nuclear factor kappa-B ligand; TNF-related activation-induced cytokine; anti-CD254 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CD254 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CD254. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CD254 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of CD254 staining in human lymph node formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CD254 expression in Jurkat (A), Raji (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CD254 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507595
NCBI GeneID	8600
NCBI Accession #	NP_003692.1
NCBI GenBank Nucleotide #	NM_003701.3
UniProt Accession #	O14788; O14723; Q96Q17; Q9P2Q3
Molecular Weight	35,478 Da
NCBI Official Full Name	tumor necrosis factor ligand superfamily member 11 isoform 1
NCBI Official Synonym Full Names	tumor necrosis factor (ligand) superfamily, member 11
NCBI Official Symbol	TNFSF11
NCBI Official Synonym Symbols	ODF; OPGL; sOdf; CD254; OPTB2; RANKL; TRANCE; hRANKL2
NCBI Protein Information	tumor necrosis factor ligand superfamily member 11; osteoprotegerin ligand; osteoclast differentiation factor; TNF-related activation-induced cytokine; receptor activator of nuclear factor kappa B ligand; receptor activator of nuclear factor kappa-B ligand
UniProt Protein Name	Tumor necrosis factor ligand superfamily member 11
UniProt Gene Name	TNFSF11
UniProt Synonym Gene Names	OPGL; RANKL; TRANCE; ODF; OPGL; RANKL; TRANCE
UniProt Entry Name	TNF11_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd254-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220489>