

Rabbit Collagen 6 alpha 2 Polyclonal Antibody | anti-COL6A2 antibody

Rabbit anti-Collagen 6 alpha 2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220873	Rabbit	Human, Monkey	Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Collagen 6 alpha 2; Polyclonal Antibody; Rabbit anti-Collagen 6 alpha 2 Antibody; Collagen alpha-2(VI) chain; anti-COL6A2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Monkey
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-COL6A2 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Collagen 6 alpha 2. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Collagen 6 alpha 2 staining in HT29 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Collagen 6 alpha 2 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Collagen 6 alpha 2 expression in HepG2 (A), HT29 (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-COL6A2 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	115527062
NCBI GeneID	1292
NCBI Accession #	NP_001840.3
NCBI GenBank Nucleotide #	NM_001849.3
UniProt Accession #	P12110; Q13909; Q13910; Q13911; Q14048; Q14049; Q16259; Q16597; Q6P0Q1; Q9UML3; Q9Y4S8
Molecular Weight	108,579 Da
NCBI Official Full Name	collagen alpha-2(VI) chain isoform 2C2
NCBI Official Synonym Full Names	collagen, type VI, alpha 2
NCBI Official Symbol	COL6A2
NCBI Official Synonym Symbols	PP3610
NCBI Protein Information	collagen alpha-2(VI) chain; collagen alpha-2(VI) chain; collagen VI, alpha-2 polypeptide; human mRNA for collagen VI alpha-2 C-terminal globular domain
UniProt Protein Name	Collagen alpha-2(VI) chain
UniProt Gene Name	COL6A2
UniProt Entry Name	CO6A2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/collagen-6-alpha-2-antibody-immunoprecipitation-immunofluorescence-immunohistochemistry-western-blot-elisa/220873>