

Rabbit NAP1 Polyclonal Antibody | anti-NAP1 antibody

Rabbit anti-NAP1 Antibody

CATALOG # AAA220932	HOST Rabbit	REACTIVITY Human, Monkey	APPLICATIONS Immunofluorescence, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	------------------------------------	---

PRODUCT SYNONYMS

NAP1; Polyclonal Antibody; Rabbit anti-NAP1 Antibody; TBKBP2; 5-azacytidine-induced protein 2; NF-kappa-B-activating kinase-associated protein 1; Nak-associated protein 1; TILP; anti-NAP1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Monkey
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-NAP1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human NAP1. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of NAP1 staining in THP1 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of NAP1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of NAP1 expression in HeLa (A), THP1 (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-NAP1 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	11968003
NCBI GeneID	64343
NCBI Accession #	NP_071906.1
NCBI GenBank Nucleotide #	NM_022461.4
UniProt Accession #	Q9H6S1; Q86W99; Q9BQF1; A8K3M2
Molecular Weight	44,935 Da
NCBI Official Full Name	5-azacytidine-induced protein 2 isoform a
NCBI Official Synonym Full Names	5-azacytidine induced 2
NCBI Official Symbol	AZI2
NCBI Official Synonym Symbols	AZ2; NAP1; TILP
NCBI Protein Information	5-azacytidine-induced protein 2; NAK-associated protein 1; 5-azacytidine induced gene 2; NF-kappa-B-activating kinase-associated protein 1
UniProt Protein Name	5-azacytidine-induced protein 2
UniProt Gene Name	AZI2
UniProt Synonym Gene Names	NAP1; Nak-associated protein 1
UniProt Entry Name	AZI2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/nap1-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220932>