

Rabbit CD172a Polyclonal Antibody | anti-CD172a antibody

Rabbit anti-CD172a Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220942	Rabbit	Human, Monkey	Immunoprecipitation, Immunofluorescence, Western Blot


PRODUCT SYNONYMS

CD172a; Polyclonal Antibody; Rabbit anti-CD172a Antibody; BIT; MFR; MYD1; PTPNS1; SHPS1; SIRP; Tyrosine-protein phosphatase non-receptor type substrate 1; SHP substrate 1; SHPS-1; Brain Ig-like molecule with tyrosine-based activation motifs; Bit; CD172 antigen-like family member A; Inhibitory receptor SHPS-1; Macrophage fusion receptor; MyD-1 antigen; Signal-regulatory protein alpha-1; Sirp-alpha-1; Signal-regulatory protein alpha-2; Sirp-alpha-2; Signal-regulatory protein alpha-3; Sirp-alpha-3; p84; anti-CD172a antibody


PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Monkey
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CD172a antibody	IP (Immunoprecipitation), IC (Immunochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CD172a. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES


IF (Immunofluorescence)

Immunofluorescent analysis of CD172a staining in THP1 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).


IHC (Immunohistochemistry)



WB (Western Blot)

Western blot analysis of CD172a expression in THP1 (A), AML193 (B), HL60 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CD172a antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	91105764
NCBI GeneID	140885
NCBI Accession #	NP_001035111.1
NCBI GenBank Nucleotide #	NM_001040022.1
UniProt Accession #	P78324; O00683; O43799; Q8N517; Q8TAL8; Q9H0Z2; Q9UDX2; Q9UIJ6; Q9Y4U9; A2A2E1; A8K411; B2R6C3
Molecular Weight	54,967 Da
NCBI Official Full Name	tyrosine-protein phosphatase non-receptor type substrate 1
NCBI Official Synonym Full Names	signal-regulatory protein alpha
NCBI Official Symbol	SIRPA
NCBI Official Synonym Symbols	BIT; MFR; P84; SIRP; MYD-1; SHPS1; CD172A; PTPNS1
NCBI Protein Information	tyrosine-protein phosphatase non-receptor type substrate 1; myd-1 antigen; inhibitory receptor SHPS-1; macrophage fusion receptor; CD172 antigen-like family member A; tyrosine phosphatase SHP substrate 1; brain-immunoglobulin-like molecule with tyrosine-b
UniProt Protein Name	Tyrosine-protein phosphatase non-receptor type substrate 1
UniProt Gene Name	SIRPA
UniProt Synonym Gene Names	BIT; MFR; MYD1; PTPNS1; SHPS1; SIRP; SHP substrate 1; SHPS-1; Bit; Sirp-alpha-1; Sirp-alpha-2
UniProt Entry Name	SHPS1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd172a-antibody-immunoprecipitation-immunochemistry-immunofluorescence-western-blot-elisa/220942>