

Rabbit anti-Human, Mouse Galectin 8 Polyclonal Antibody | anti-LGALS8 antibody

Rabbit anti-Galectin 8 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220950	Rabbit	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Galectin 8; Polyclonal Antibody; Rabbit anti-Galectin 8 Antibody; Galectin-8; Gal-8; Po66 carbohydrate-binding protein; Po66-CBP; Prostate carcinoma tumor antigen 1; PCTA-1; anti-LGALS8 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-LGALS8 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Galectin 8. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Galectin 8 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Galectin 8 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Galectin 8 expression in HepG2 (A), NIH3T3 (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-LGALS8 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	42544185
NCBI GeneID	3964
NCBI Accession #	NP_006490.3
NCBI GenBank Nucleotide #	NM_006499.4
UniProt Accession #	O00214; Q9JL15
Molecular Weight	35,808 Da
NCBI Official Full Name	galectin-8 isoform a
NCBI Official Synonym Full Names	lectin, galactoside-binding, soluble, 8
NCBI Official Symbol	LGALS8
NCBI Official Synonym Symbols	Gal-8; PCTA1; PCTA-1; Po66-CBP
NCBI Protein Information	galectin-8; galectin 8; galectin-8g; Po66 carbohydrate binding protein; po66 carbohydrate-binding protein; prostate carcinoma tumor antigen 1
UniProt Protein Name	Galectin-8
UniProt Gene Name	LGALS8
UniProt Synonym Gene Names	Gal-8; Po66-CBP; PCTA-1
UniProt Entry Name	LEG8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/galectin-8-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220950>