

Rabbit Androgen Receptor Polyclonal Antibody | anti-AR antibody

Rabbit anti-Androgen Receptor (pS213) Antibody

CATALOG # AAA220984	HOST Rabbit	REACTIVITY Human, Monkey	APPLICATIONS Immunofluorescence, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	------------------------------------	---

PRODUCT SYNONYMS

Androgen Receptor; Polyclonal Antibody; Rabbit anti-Androgen Receptor (pS213) Antibody; DHTR; NR3C4; Androgen receptor; Dihydrotestosterone receptor; Nuclear receptor subfamily 3 group C member 4; anti-AR antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Monkey
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-AR antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Androgen Receptor. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Androgen Receptor (pS213) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Androgen Receptor (pS213) staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Androgen Receptor (pS213) expression in HepG2 (A), HeLa (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-AR antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	21322252
NCBI GeneID	367
NCBI Accession #	NP_000035.2
NCBI GenBank Nucleotide #	NM_000044.3
UniProt Accession #	P10275; Q9UD95; A2RUN2; B1AKD7
Molecular Weight	98,989 Da
NCBI Official Full Name	androgen receptor isoform 1
NCBI Official Synonym Full Names	androgen receptor
NCBI Official Symbol	AR
NCBI Official Synonym Symbols	KD; AIS; TFM; DHTR; SBMA; HYS P1; NR3C4; SMAX1; HUMARA
NCBI Protein Information	androgen receptor; dihydrotestosterone receptor; androgen nuclear receptor variant 2; nuclear receptor subfamily 3 group C member 4
UniProt Protein Name	Androgen receptor
UniProt Gene Name	AR
UniProt Synonym Gene Names	DHTR; NR3C4
UniProt Entry Name	ANDR_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/androgen-receptor-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220984>