

Rabbit ICK Polyclonal Antibody | anti-ICK antibody

Rabbit anti-ICK Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA221023	Rabbit	Human, Bovine	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

ICK; Polyclonal Antibody; Rabbit anti-ICK Antibody; KIAA0936; Serine/threonine-protein kinase ICK; Intestinal cell kinase; hICK; Laryngeal cancer kinase 2; LCK2; MAK-related kinase; MRK; anti-ICK antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Bovine
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-ICK antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ICK. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of ICK staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of ICK staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of ICK expression in HeLa (A), HEK293T (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-ICK antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	7662388
NCBI GeneID	22858
NCBI Accession #	NP_055735.1
NCBI GenBank Nucleotide #	NM_014920.3
UniProt Accession #	Q9UPZ9; O75985; Q5THL2; Q8IYH8; Q9BX17; Q9NYX3; A7MD41
Molecular Weight	71,427 Da
NCBI Official Full Name	serine/threonine-protein kinase ICK
NCBI Official Synonym Full Names	intestinal cell (MAK-like) kinase
NCBI Official Symbol	ICK
NCBI Official Synonym Symbols	ECO; MRK; LCK2
NCBI Protein Information	serine/threonine-protein kinase ICK; hICK; MAK-related kinase; laryngeal cancer kinase 2; serine/threonine protein kinase
UniProt Protein Name	Serine/threonine-protein kinase ICK
UniProt Gene Name	ICK
UniProt Synonym Gene Names	KIAA0936; hICK; LCK2; MRK
UniProt Entry Name	ICK_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/ick-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/221023>