

Rabbit anti-Human, Mouse DAP Kinase 1 Polyclonal Antibody | anti-DAP antibody

Rabbit anti-DAP Kinase 1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA221054	Rabbit	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

DAP Kinase 1; Polyclonal Antibody; Rabbit anti-DAP Kinase 1 Antibody; DAPK; Death-associated protein kinase 1; DAP kinase 1; anti-DAP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-DAP antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human DAP Kinase 1. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of DAP Kinase 1 staining in K562 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of DAP Kinase 1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of DAP Kinase 1 expression in K562 (A), mouse brain (B), NIH3T3 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-DAP antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	89363047
NCBI GeneID	1612
NCBI Accession #	NP_004929.2
NCBI GenBank Nucleotide #	NM_004938.2
UniProt Accession #	P53355; Q80YE7
Molecular Weight	1430
NCBI Official Full Name	death-associated protein kinase 1
NCBI Official Synonym Full Names	death-associated protein kinase 1
NCBI Official Symbol	DAPK1
NCBI Official Synonym Symbols	DAPK
NCBI Protein Information	death-associated protein kinase 1; DAP kinase 1
UniProt Protein Name	Death-associated protein kinase 1
UniProt Gene Name	DAPK1
UniProt Synonym Gene Names	DAPK; DAP kinase 1
UniProt Entry Name	DAPK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://stage.aaabiotech.com/polyclonal/dap-kinase-1-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/221054>