

Rabbit anti-Human, Rat APOBEC3G Polyclonal Antibody | anti-APOBEC3G antibody

Rabbit anti-APOBEC3G Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA221059	Rabbit	Human, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

APOBEC3G; Polyclonal Antibody; Rabbit anti-APOBEC3G Antibody; DNA dC->dU-editing enzyme APOBEC-3G; APOBEC-related cytidine deaminase; APOBEC-related protein; ARCD; APOBEC-related protein 9; ARP-9; CEM-15; CEM15; Deoxycytidine deaminase; A3G; anti-APOBEC3G antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-APOBEC3G antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Recombinant full length protein of human APOBEC3G
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of APOBEC3G staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of APOBEC3G staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of APOBEC3G expression in Jurkat (A) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-APOBEC3G antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	44887683
NCBI GeneID	60489
UniProt Accession #	Q9HC16; Q45F02; Q5TF77; Q7Z2N1; Q7Z2N4; Q9H9H8; B2RDR9
Molecular Weight	384
NCBI Official Full Name	DNA dC->gt
NCBI Official Synonym Full Names	apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3G
NCBI Official Symbol	APOBEC3G
NCBI Official Synonym Symbols	A3G; ARCD; ARP9; ARP-9; CEM15; CEM-15; MDS019; bK150C2.7; dJ494G10.1
NCBI Protein Information	DNA dC->dU-editing enzyme APOBEC-3G; deoxycytidine deaminase; APOBEC-related protein 9; DNA dC->dU editing enzyme; phorbolin-like protein MDS019; APOBEC-related cytidine deaminase; apolipoprotein B mRNA editing enzyme cytidine deaminase; apolipoprot
UniProt Protein Name	DNA dC->dU-editing enzyme APOBEC-3G
UniProt Gene Name	APOBEC3G
UniProt Synonym Gene Names	APOBEC-related protein; ARCD; ARP-9; CEM15; A3G
UniProt Entry Name	ABC3G_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/apobec3g-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/221059>