

Rabbit Collagen 1 alpha 2 Polyclonal Antibody | anti-COL1A2 antibody

Rabbit anti-Collagen 1 alpha 2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222459	Rabbit	Human, Dog, Rabbit	Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Collagen 1 alpha 2; Polyclonal Antibody; Rabbit anti-Collagen 1 alpha 2 Antibody; Collagen alpha-2(I) chain; Alpha-2 type I collagen; anti-COL1A2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Dog, Rabbit
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-COL1A2 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Collagen 1 alpha 2. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Collagen 1 alpha 2 staining in A431 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of Collagen 1 alpha 2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of Collagen 1 alpha 2 expression in A431 (A), HEK293T (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-COL1A2 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	48762934
NCBI GeneID	1278
NCBI Accession #	NP_000080.2
NCBI GenBank Nucleotide #	NM_000089.3
UniProt Accession #	P08123; P02464; Q13897; Q13997; Q13998; Q14038; Q14057; Q15177; Q15947; Q16480; Q16511; Q7Z5S6
Molecular Weight	129,314 Da
NCBI Official Full Name	collagen alpha-2(I) chain
NCBI Official Synonym Full Names	collagen, type I, alpha 2
NCBI Official Symbol	COL1A2
NCBI Official Synonym Symbols	OI4
NCBI Protein Information	collagen alpha-2(I) chain; collagen alpha-2(I) chain; type I procollagen; alpha 2(I)-collagen; alpha-2 type I collagen; collagen I, alpha-2 polypeptide; collagen of skin, tendon and bone, alpha-2 chain
UniProt Protein Name	Collagen alpha-2(I) chain
UniProt Gene Name	COL1A2
UniProt Entry Name	CO1A2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.