

Rabbit A20 Polyclonal Antibody | anti-A20 antibody

Rabbit anti-A20 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222524	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

A20; Polyclonal Antibody; Rabbit anti-A20 Antibody; OTUD7C; Tumor necrosis factor alpha-induced protein 3; TNF alpha-induced protein 3; OTU domain-containing protein 7C; Putative DNA-binding protein A20; Zinc finger protein A20; anti-A20 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-A20 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human A20. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of A20 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of A20 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of A20 expression in HeLa (A), Daudi (B), RAW264.7 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-A20 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	395393995
NCBI GeneID	7128
NCBI Accession #	NP_001257436.1
NCBI GenBank Nucleotide #	NM_001270507.1
UniProt Accession #	P21580; Q60769
Molecular Weight	89,614 Da
NCBI Official Full Name	tumor necrosis factor alpha-induced protein 3
NCBI Official Synonym Full Names	tumor necrosis factor, alpha-induced protein 3
NCBI Official Symbol	TNFAIP3
NCBI Official Synonym Symbols	A20; OTUD7C; TNFA1P2
NCBI Protein Information	tumor necrosis factor alpha-induced protein 3; zinc finger protein A20; TNF alpha-induced protein 3; OTU domain-containing protein 7C; putative DNA-binding protein A20; tumor necrosis factor inducible protein A20
UniProt Protein Name	Tumor necrosis factor alpha-induced protein 3
UniProt Gene Name	TNFAIP3
UniProt Synonym Gene Names	OTUD7C; TNF alpha-induced protein 3
UniProt Entry Name	TNAP3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/a20-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222524>