

Rabbit AS160 Polyclonal Antibody | anti-AS160 antibody

Rabbit anti-AS160 (pT642) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222541	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

AS160; Polyclonal Antibody; Rabbit anti-AS160 (pT642) Antibody; KIAA0603; TBC1 domain family member 4; Akt substrate of 160 kDa; anti-AS160 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-AS160 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human AS160. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of AS160 (pT642) staining in Raw264.7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of AS160 (pT642) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of AS160 (pT642) expression in LO2 insulin-treated (A), Raw264.7 insulin-treated (B), PC12 insulin-treated (C), NIH3T3 UV-treated (D) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-AS160 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	114688046
NCBI GeneID	9882
NCBI Accession #	NP_055647.2
NCBI GenBank Nucleotide #	NM_014832.2
UniProt Accession #	O60343; Q8BYJ6
Molecular Weight	146,563 Da
NCBI Official Full Name	TBC1 domain family member 4
NCBI Official Synonym Full Names	TBC1 domain family, member 4
NCBI Official Symbol	TBC1D4
NCBI Official Synonym Symbols	AS160
NCBI Protein Information	TBC1 domain family member 4; akt substrate of 160 kDa; Acrg embryonic lethality minimal region ortholog; TBC (Tre-2, BUB2, CDC16) domain-containing protein
UniProt Protein Name	TBC1 domain family member 4
UniProt Gene Name	TBC1D4
UniProt Synonym Gene Names	AS160; KIAA0603; AS160
UniProt Entry Name	TBCD4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/as160-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222541>