

Rabbit Cyclin E1 Polyclonal Antibody | anti-CCNE1 antibody

Rabbit anti-Cyclin E1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222583	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Cyclin E1; Polyclonal Antibody; Rabbit anti-Cyclin E1 Antibody; CCNE; G1/S-specific cyclin-E1; anti-CCNE1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CCNE1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cyclin E1. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Cyclin E1 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Cyclin E1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of Cyclin E1 expression in MCF7 (A), mouse liver (B), rat kidney (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CCNE1 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	17318559
NCBI GeneID	898
NCBI Accession #	NP_001229.1
NCBI GenBank Nucleotide #	NM_001238.2
UniProt Accession #	P24864; Q61457; P39949
Molecular Weight	47,077 Da
NCBI Official Full Name	G1/S-specific cyclin-E1
NCBI Official Synonym Full Names	cyclin E1
NCBI Official Symbol	CCNE1
NCBI Official Synonym Symbols	CCNE
NCBI Protein Information	G1/S-specific cyclin-E1; cyclin Es; cyclin Et
UniProt Protein Name	G1/S-specific cyclin-E1
UniProt Gene Name	CCNE1
UniProt Synonym Gene Names	CCNE
UniProt Entry Name	CCNE1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.