

Rabbit GADD153 Polyclonal Antibody | anti-GADD153 antibody

Rabbit anti-GADD153 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222587	Rabbit	Human, Monkey, Pig	Immunoprecipitation, Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

GADD153; Polyclonal Antibody; Rabbit anti-GADD153 Antibody; CHOP; CHOP10; DNA damage-inducible transcript 3 protein; DDIT-3; C/EBP zeta; C/EBP-homologous protein; C/EBP-homologous protein 10; CHOP-10; CCAAT/enhancer-binding protein homologous protein; Growth arrest and DNA damage-inducible protein GADD153; anti-GADD153 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Monkey, Pig
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-GADD153 antibody	IP (Immunoprecipitation), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human GADD153. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of GADD153 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of GADD153 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of GADD153 expression in Jurkat (A), HeLa (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-GADD153 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	304282234
NCBI GeneID	1649
NCBI Accession #	NP_001181986.1
NCBI GenBank Nucleotide #	NM_001195057.1
UniProt Accession #	P35638
Molecular Weight	19,175 Da
NCBI Official Full Name	DNA damage-inducible transcript 3 protein isoform 2
NCBI Official Synonym Full Names	DNA-damage-inducible transcript 3
NCBI Official Symbol	DDIT3
NCBI Official Synonym Symbols	CHOP; CEBPZ; CHOP10; CHOP-10; GADD153
NCBI Protein Information	DNA damage-inducible transcript 3 protein; DDIT-3; C/EBP zeta; c/EBP-homologous protein 10; CCAAT/enhancer-binding protein homologous protein; growth arrest and DNA damage-inducible protein GADD153
UniProt Protein Name	DNA damage-inducible transcript 3 protein
UniProt Gene Name	DDIT3
UniProt Synonym Gene Names	CHOP; CHOP10; GADD153; DDIT-3; CHOP; CHOP-10
UniProt Entry Name	DDIT3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/gadd153-antibody-immunoprecipitation-immunofluorescence-immunohistochemistry-western-blot-elisa/222587>