

Rabbit TRAF3IP1 Polyclonal Antibody | anti-TRAF3IP1 antibody

Rabbit anti-TRAF3IP1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222610	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

TRAF3IP1; Polyclonal Antibody; Rabbit anti-TRAF3IP1 Antibody; MIPT3; TRAF3-interacting protein 1; Interleukin-13 receptor alpha 1-binding protein 1; Microtubule-interacting protein associated with TRAF3; MIP-T3; anti-TRAF3IP1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-TRAF3IP1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TRAF3IP1. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of TRAF3IP1 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of TRAF3IP1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of TRAF3IP1 expression in U87 (A), HeLa (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-TRAF3IP1 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	213385278
NCBI GeneID	26146
NCBI Accession #	NP_001132962.1
NCBI GenBank Nucleotide #	NM_001139490.1
UniProt Accession #	Q8TDR0; Q149C2; Q5XIN3
Molecular Weight	78,632 Da
NCBI Official Full Name	TRAF3-interacting protein 1 isoform 2
NCBI Official Synonym Full Names	TNF receptor-associated factor 3 interacting protein 1
NCBI Official Symbol	TRAF3IP1
NCBI Official Synonym Symbols	MIPT3; MIP-T3
NCBI Protein Information	TRAF3-interacting protein 1; interleukin 13 receptor alpha 1-binding protein-1; interleukin-13 receptor alpha 1-binding protein 1; microtubule-interacting protein associated with TRAF3; microtubule interacting protein that associates with TRAF3; microtubu
UniProt Protein Name	TRAF3-interacting protein 1
UniProt Gene Name	TRAF3IP1
UniProt Synonym Gene Names	MIPT3; MIP-T3
UniProt Entry Name	MIPT3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/traf3ip1-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222610>