

Rabbit Histone Deacetylase 8 Polyclonal Antibody | anti-HDAC8 antibody

Rabbit anti-Histone Deacetylase 8 (pS39) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222612	Rabbit	Human, Mouse, Rat, Monkey	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Histone Deacetylase 8; Polyclonal Antibody; Rabbit anti-Histone Deacetylase 8 (pS39) Antibody; HDAC1; Histone deacetylase 8; HD8; anti-HDAC8 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Monkey
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-HDAC8 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Histone Deacetylase 8 (pS39). The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Histone Deacetylase 8 (pS39) staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Histone Deacetylase 8 (pS39) staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Histone Deacetylase 8 (pS39) expression in A549 (A), NIH3T3 (B), rat kidney (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-HDAC8 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	29839394
NCBI GeneID	55869
UniProt Accession #	Q9BY41; Q8VH37; B1WC68
Molecular Weight	377
NCBI Official Full Name	Histone deacetylase 8
NCBI Official Synonym Full Names	histone deacetylase 8
NCBI Official Symbol	HDAC8
NCBI Official Synonym Symbols	HD8; WTS; RPD3; CDA07; CDLS5; MRXS6; HDACL1
NCBI Protein Information	histone deacetylase 8; histone deacetylase-like 1
UniProt Protein Name	Histone deacetylase 8
UniProt Gene Name	HDAC8
UniProt Synonym Gene Names	HDACL1; HD8
UniProt Entry Name	HDAC8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/histone-deacetylase-8-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222612>