

Rabbit GAB1 Polyclonal Antibody | anti-GAB1 antibody

Rabbit anti-GAB1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222676	Rabbit	Human, Mouse, Bovine	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

GAB1; Polyclonal Antibody; Rabbit anti-GAB1 Antibody; GRB2-associated-binding protein 1; GRB2-associated binder 1; Growth factor receptor bound protein 2-associated protein 1; anti-GAB1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Bovine
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-GAB1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human GAB1. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of GAB1 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of GAB1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of GAB1 expression in HepG2 (A), Raw264.7 (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-GAB1 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	46370073
NCBI GeneID	2549
NCBI Accession #	NP_002030.2
NCBI GenBank Nucleotide #	NM_002039.3
UniProt Accession #	Q13480; Q9QYY0
Molecular Weight	76,616 Da
NCBI Official Full Name	GRB2-associated-binding protein 1 isoform b
NCBI Official Synonym Full Names	GRB2-associated binding protein 1
NCBI Official Symbol	GAB1
NCBI Protein Information	GRB2-associated-binding protein 1; GRB2-associated binder 1; growth factor receptor bound protein 2-associated protein 1
UniProt Protein Name	GRB2-associated-binding protein 1
UniProt Gene Name	GAB1
UniProt Entry Name	GAB1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/gab1-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222676>