

Rabbit TRK A Polyclonal Antibody | anti-TRKA antibody

Rabbit anti-TRK A (pY496) Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222719	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

TRK A; Polyclonal Antibody; Rabbit anti-TRK A (pY496) Antibody; MTC; TRK; TRKA; High affinity nerve growth factor receptor; Neurotrophic tyrosine kinase receptor type 1; TRK1-transforming tyrosine kinase protein; Tropomyosin-related kinase A; Tyrosine kinase receptor; Tyrosine kinase receptor A; Trk-A; gp140trk; p140-TrkA; anti-TRKA antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-TRKA antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TRK A. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of TRK A (pY496) staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of TRK A (pY496) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of TRK A (pY496) expression in mouse brain (A), rat brain (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-TRKA antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	56118210
NCBI GeneID	4914
NCBI Accession #	NP_001007793.1
NCBI GenBank Nucleotide #	NM_001007792.1
UniProt Accession #	P04629; Q3UFB7; P35739
Molecular Weight	87,497 Da
NCBI Official Full Name	high affinity nerve growth factor receptor isoform 3
NCBI Official Synonym Full Names	neurotrophic tyrosine kinase, receptor, type 1
NCBI Official Symbol	NTRK1
NCBI Official Synonym Symbols	MTC; TRK; TRK1; TRKA; Trk-A; p140-TrkA
NCBI Protein Information	high affinity nerve growth factor receptor; gp140trk; Oncogene TRK; tyrosine kinase receptor A; tropomyosin-related kinase A; TRK1-transforming tyrosine kinase protein
UniProt Protein Name	High affinity nerve growth factor receptor
UniProt Gene Name	NTRK1
UniProt Synonym Gene Names	MTC; TRK; TRKA; Trk-A
UniProt Entry Name	NTRK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/trk-a-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222719>